



AGRICULTURE

NOTICE: Return or ranew all Library Materials! The Minimum Fee for each Lost Book is \$50.00.

The person charging this material is responsible for its return to the library from which it was withdrawn on or before the **Latest Date** stamped below.

Theft, mutilation, and underlining of books are reasons for disciplinary action and may result in dismissal from the University. To renew call Telephone Center, 333-8400

UNIVERSITY OF ILLINOIS LIBRARY AT URBANA-CHAMPAIGN

APR 2 1 1996 L161-O-1096







1947 Illinois HYBRID CORN **TESTS**

Bulletin 527

UNIVERSITY OF ILLINOIS

AGRICULTURAL EXPERIMENT STATION in cooperation with ILLINOIS STATE NATURAL HISTORY SURVEY . . . February, 1948

Table 1. — GENERAL INFORMATION: Illinois Cooperative Hybrid Corn Tests, 1947

Field	County and location	Num- ber of	Date planted	Date harvested		rage yield	Average mois- ture in	Average erect
	in state	ntrie		nai vesteu	Total	Sound	grain	plants
					bu.	bu.	perct.	perct.
Woodstock			May 31	Nov. 4	42.4	41.5	28.4	81.6
Galesburg			May 24	Nov. 10, 11	73.0	71.6	22.3	63.1
Sheldon			June 5	Nov. 12	71.1	70.2	24.6	82.8
Sullivan	Moultrie (SC)	81	May 24	Oct. 31	63.8	62.4	20.6	97.5
Alhambra Robbs (Dixon Sp.)	Madison (S)	81	June 11	Dec. 11, 12	48.5	47.4	23.7	55.3
Bottomland	Pope (Ex.S)	60	June 16	Nov. 24, 25	41.7	40.9	26.5	80.4
Upland			May 28	Nov. 25 .	34.0	33.5	21.6	78.7

COOPERATORS: Earl Hughes, McHenry county; Earl and Webster Gehring, Knox county; John B. Rice, Iroquois county; R. B. Vandeveer, Farm Manager, Illinois Masonic Home Farm, Moultric county. The Alhambra field in Madison county is managed by the Illinois Station. The Pope county fields at Robbs are parts of the Dixon Springs Experiment Station of which R. J. Webb is superintendent and J. M. Lewis is assistant superintendent.

Most of the hybrids selected for testing are extensively grown in the state. Some experimental hybrids were included because they had shown promise for commercial production in preliminary tests. A few hybrids were put in the test mainly to meet the field-performance requirement for certification.

Open-pedigree hybrids from commercial seed producers were entered in the 1947 test. The name of the producer of each lot of open-pedigree hybrid seed is included in the tables.

Soil characteristics of fields. The test fields were medium to high in productivity, and each represents a soil type common to the region where it is located. Each field was selected carefully for uniformity in soil type, productivity, and drainage. The Alhambra field contained a number of "slick spots" and was the most variable in productivity.

In 1947 the northern Illinois test was moved to McHenry county. The other tests were conducted on the same farms as in 1946. The approximate locations of the test fields are shown on the map on the inside front cover. General information on soil characteristics and soil management is given in Table 2.

Field-plot design. A 9 x 9 randomized, lattice-square field-plot design with 5 replications was used on the Woodstock, Galesburg, Sheldon, Sullivan, and Alhambra fields. Controlled, randomized block designs with 6 replications were used on the Dixon Springs bottomland field and 5 on the upland field.

Method of planting. All test fields were planted by hand

Table 2.—TESTING FIELDS: Soil Characteristics and Management Practices

Soil type	Lime require- ment	Available phosphorus	Available potassium	Previous crops and soil management
	NC	RTHERN: V	Woodstock	
Proctor silt loam, flat phase	tons	Medium	Low to medium	Oats 1944; alfalfa 1945-1946; 3 tons lime and 240 pounds 3-12-12 1944; 500 pounds rock phos- phate 1942.
	WEST N	ORTH-CENT	RAL: Gales	burg
Muscatine silt loam	3	Medium	Very high	Corn 1944; oats 1945; red clover hog pasture 1946; lime and rock phosphate have been applied.
	EAST N	ORTH-CENT	TRAL: Shel	don
Drummer clay loam	0	High	Very high	Corn 1942; oats 1943; alfalfa 1944- 1946; 2 tons lime 1943; ½ ton rock phosphate 1938; 125 pounds potash 1946.
	sou	TH-CENTRA	L: Sullivan	
Flanagan silt loam	2	Medium	High	Alfalfa 1941-1943; corn 1944; alfalfa-timothy pasture 1945-1946; 2 tons lime 1946.
	S	OUTHERN:	Alhambra	
Putnam silt loam	0	High	High	Wheat 1941; corn 1942; oats 1943; soybeans 1944; wheat 1945; red clover 1946; 500 pounds potash broadcast 1947 before planting.
EXT	REME SO	OUTHERN: 1	Robbs (Dixo	n Springs)
Upland field: Ava silt loam.	trace	Very low	High	Timothy-sweet clover sod past 15 years.
Bottomland field: Bonnie silt loam		Very low	Very high	Corn 1944; soybeans 1945; corn 1946; no soil treatment.

R. S. SMITH, Chief in Soil Physics and Soil Survey, and HERMAN WASCHER, Associate Chief in Soil Survey, have approved the soil-type designations, uniformity, and physical characteristics of the above fields.

on land prepared in the regular way for corn. Each plot consisted of 2 rows 10 hills long, except at Alhambra where the plots were only 9 hills long. Three kernels were dropped in each hill except on the fields at Dixon Springs, where only 2 kernels were planted.

Data from all plots were included in the results. The only correction for imperfect stand was the following adjustment for missing hills:

Ear weight in field
$$\times \left(1 + \frac{\text{missing hills}}{\text{hills present}} \times .6\right) =$$
 adjusted ear weight.

WEATHER CONDITIONS

In general, the 1947 planting season was another extremely late and discouraging one thruout Illinois. The cool spring was so wet that lowlands and low spots in fields were repeatedly flooded. Stands were uneven and, since wet fields prevented timely cultivation, were more weedy than usual.

Wet weather made the planting of test fields later than usual. All fields except the Sheldon field were planted in good seedbeds which had ample moisture and were planted under conditions favorable for growth. The test area on the Sheldon field was planted five days after the rest of the field. A heavy rain which fell at sometime during the five-day period made the originally excellent seedbed very compact.

Good stands were obtained on the Woodstock, Galesburg, Sullivan, and Dixon Springs bottomland fields. Stands were variable on the Sheldon, Alhambra, and Dixon Springs upland fields.

During June corn made good growth despite the cool, wet weather. Corn suffered during late July and August from the excessive heat and drouth. Late-planted corn suffered, particularly when its early growth was extremely rapid. Heat caused stunting and firing; this condition was more general in the southern half of the state and the damage most serious there.

Serious fall damage from frost had threatened the late-planted Illinois corn crop since planting time. The fall season, however, was unusually warm, killing frosts generally holding off until early November. The exception was the extreme northern part of the state, where a killing frost occurred during the last week of September. The Woodstock field was in this area but the damage was slight. Clear, warm weather during September and October hastened drying thruout the state. Practically all the corn matured without serious damage and the expected soft corn drying, handling, and storage problems did not materialize.

INSECT PESTS

European corn borer. In northern Illinois in 1947, corn losses from European corn borer, *Pyrausta nubilalis* (Hbn.), were material. Hybrid test fields at Woodstock and Sheldon were in this

area. Records of stalk breakage in these fields at harvest time are shown in Table 5 (page 536) and Table 10 (page 543).

The percentage of plants broken below the ear because of borer attack and the average breakage for the fields show that the attack varied in intensity at the two fields. At Woodstock, plants broken below the ear ranged from less than one percent to over 23 percent; the average was 8.4 percent. The difference necessary for significance in hybrids is 8 percent (Table 5). Many hybrids appear to be very similar in their response to borer attack. A three-year average for 19 hybrids shows that 2.8 to 9.8 percent of the plants were broken below the ear (Table 5). The difference between hybrids in this three-year average is not significant.

The percentage of plants broken below the ear at Sheldon range from less than 1 percent to 10.9 percent. In this field, having little breakage, differences between hybrids in the test is not significant. A three-year average of 26 entries shows that the percentage of plants broken below the ear ranged from 14.6 to 25.5. The average for all entries was 19.7 percent. None is significantly better than any other.

Southern corn rootworm. The hybrid test field at Galesburg was attacked by corn rootworms, especially southern corn rootworm, Diabrotica duodecimpunctata (F.). The attack damaged root systems and caused lodging, which was greatly increased by a windstorm. Forty-five of the hybrids in the field showed better than average resistance to lodging. Total lodging, however, ranged widely — from 2.2 to 74.3 percent, the average being 33.6 percent (Table 8, page 540). Because the windstorm caused a big difference between portions of the field, a difference of 33.6 percent is necessary for significance (Table 8). Figures for lodging are available for the years 1943 and 1947; average figures for 22 hybrids grown on this field during those years are given in Table 7 (page 539). Differences between hybrids included in these averages are not significant.

DISEASE DAMAGE1

Stalk rot diseases. From surveys in 37 scattered counties, direct damage from stalk rot was estimated at 5 percent. In some fields the stalks were badly broken, the breakage resulting in indirect losses since the pickers did not get all the ears. The amount of damage varied a good deal from place to place. Of the stalks examined, 33 percent were rotted by *Diplodia zeae* and 25 percent by *Gibberella zeae*. In northern Illinois, Gibberella was more prevalent than Diplodia. In southern Illinois, 1.4 percent of the stalks were damaged by charcoal rot. *Fusarium moniliforme*, *Nigrospora oryzae*, and other organisms also caused a moderate amount of stalk rot.

The Diplodia has usually been the most serious cause of corn stalk rot in Illinois, Gibberella was the most important cause in 1946 and of considerable importance in 1947. Some inbreds that are resistant to Diplodia are not resistant to Gibberella and vice versa.

Root rot. Wet soil in the spring and compacted, poorly aerated soil during the summer were very favorable to root rot organisms. Roots were brown and plants were easily pulled up in many fields. However, a specific estimate on the amount of damage can not be made.

Smut. Loss from smut was estimated at only 0.5 percent. Damage from this disease has been moderate since 1940, in which year the loss was estimated at 4 percent.

Stewart's disease. This disease was observed especially in canning corn in Iroquois and Vermilion counties and in nonresistant garden varieties of sweet corn in central and south-central Illinois. In late summer, leaf symptoms of Stewart's disease were very common on field corn except in the northern third of the state. This infection in field corn apparently caused little direct damage, but it may have increased stalk rot susceptibility.

Ear rots. Corn kernel damage from rot in the various performance fields is given in Table 3. The dry weather of late summer and fall was not conducive to ear rot infection and the percentages are low, altho not so low as in 1946. The least ear rot damage occurred at Sheldon, the most at Alhambra where the most

^{&#}x27;Estimates of losses are based in part on survey data obtained by G. H. Boewe of the Illinois State Natural History Survey.

Table 3.—ROT DAMAGE CAUSED BY FUNGI: Average of All Entries on Six Test Fields, 1947

(Figures based on laboratory tests)

				Corn ker	nels damag	ged by ro	t	
Ran	k* Fungi causing damage	Wood- stock	Gales-	Sheldon	Sullivan	Alham- bra	Dixon S ₁	orings
	agc	Stock	burg	Sheldon	Sumvan	Dra	Bottomland	Upland
		perct.	perct.	perct.	perct.	perci.	perci.	perct.
1	Fusarium moniliforme	. 29	1.17	.32	. 40	1.24	1.55	1.47
2	Penicillium spp	.51	. 24	.09	. 13	.64	.06	.05
3	Diplodia zeae	.03	.07	.54	. 90	.02	0	0
4	Aspergillus spp	. 92	.03	0	. 13	.08	.01	.01
5	Nigrospora oryzae	.09	. 15	.09	.04	. 20	.02	.01
6	Rhizopus and Mucor	.07	.08	.03	. 13	.08	. 10	.04
7	Alternaria spp	.03	.09	.04	.12	.08	.01	0
8	Gibberella zeae	0	0	. 10	.01	.02	0	.03
	Others	.07	. 19	.06	. 15	. 12	.11	.02
	Total	2.01	2.02	1.27	2.01	2.48	1.86	1.63

^{*} Based on total damage on all fields.

lodging also occurred. In the last two years Fusarium moniliforme was the chief cause of kernel damage by rot. Penicillium jumped up to second place in 1947. At Dixon Springs Diplodia was not even found. Ear rot losses were high in 1941 and since then we have had more or less dry fall weather and low rot damage.

MEASURING PERFORMANCE

The entries in the 1947 test are listed in the tables in the order of their total yields. Two or more entries having the same total yield are given the same rating, but the one having the higher yield of sound corn is placed first. Those having the same total yield and sound yield are placed in order by percentage of erect plants.

Erect plants. The percentage of erect plants in each plot of each entry on each field was estimated at the time of harvest. The ratings for erect plants show how the percentage of erect plants for each hybrid compared with the percentage of erect plants on the field as a whole. (Each rating is obtained by dividing the percentage of erect plants for that hybrid by the percentage of erect plants on the field as a whole and multiplying by 100.)

Lodging may have been due to rootworm damage, weak or rotted roots, corn borer damage, stalk rots, or weak stalks. Stalks broken above the ear were not considered lodged.

Yield of grain. To determine shelling percentage, all the

ears from one replicate of each entry were shelled. At Dixon Springs, however, because it was not practicable to shell all the ears in a replication, the shelling percentage of all entries was assumed to be 80 percent. A sample of shelled corn was taken from the Dixon Springs plots by hand-shelling 6 ears of each entry in one replication.

From the shelled corn one sample was taken to determine the percentage of moisture at harvest¹ and to determine the percentage of damaged kernels. The percentage of damaged corn was determined according to the federal grain standards.

The total acre-yield was calculated as shelled corn containing 15.5 percent moisture, the upper limit allowable in No. 2 corn. The total yield thus obtained was adjusted according to the procedure outlined by Cochran for randomized lattice-square designs.² The total yield of sound corn was computed by deducting the amount of damaged corn from the total yield.

Each hybrid's rating for sound yield, expressed in percentage, is simply the ratio between the bushels of sound corn produced by the hybrid and the average number of bushels of sound corn produced by all the entries on the field.

Height of ear. Notes on comparative height of ear were taken at harvest time. Each plot of each entry was placed in one of the five following categories: low, mid-low (midway between low and medium), medium, mid-high (midway between medium and high), and high. Beginning with low and continuing progressively to high, these terms were assigned numerical values from 1 to 5 to permit the averaging of the plots.

Significance of yield differences. Too much confidence must not be placed in the particular ranking of a hybrid in the following tables, for chance has played a part in determining its position. Unaccountable variability in the soil and conditions on the field will cause differences in yield that are not inherent in the hybrids themselves.

The part played by chance in the 1947 tests has been calculated for total yield by the mathematical procedure known as "analysis

¹ All moisture determinations were made with a Steinlite moisture tester.

² Cochran, W. G. "Some Additional Lattice-Square Designs." *Iowa Agr. Exp. Sta. Res. Bul. 318.* May, 1943.

of variance." At the bottom of each table is stated the approximate difference which there must be between any two entries in order for them to show a true inherent difference. Unless two hybrids differ by at least this amount, there is no assurance that one hybrid is inherently higher yielding than the other.

RESULTS OF TESTS

Detailed results of the tests on six regular test fields and the two special soil-adaptation fields are given in Tables 4 to 14 on the following pages. See also Table 3 on page 531 on ear-rot damage.

Readers are urged to keep in mind these two things when comparing the performance of hybrids on any one field:

- 1. Small differences in yield do not necessarily indicate the superiority of one hybrid over another. See each table for the amount one hybrid must exceed another before it can be considered the better.
- 2. The Summary section of each table is the most important part. At least three years' results are necessary to give a reasonably reliable picture of a hybrid's ability to perform under varying seasonal conditions.

Table 4. — NORTHERN ILLINOIS: Kings and Woodstock

		Λοπο	-vield	Damaged corn in	Mois- ture in	Erect	Ratin	g for—	Compara
Rank	Entry -	Acre	-yieiu	- shelled	grain at	plants	Erect	Sound	height
		Total	Sound*	samples	harvest		plants	yield*	of ear
	SUMMAR	Y: 1	Kings	1945, 19	946; W	oodsto	ck 194	7	
		bu.	bu.	perct.	perct.	perct.	perct.	perct.	
1 F	urr 67A	68.9	68.7	. 5	27.7	79	97.4	106.0	Mediur
2 S	ieben S-450	68.4	68.1	. 4	26.0	81	99.9	105.0	Mediur
2 I	Doubet D-1	68.4	67.4	1.8	28.3	86	106.0	104.0	Mediur
4 N	lichols 5A	68.1	67.6	1.0	30.2	87	107.3	104.2	Mediur
5 I	DeKalb 609	67.9			28.7	80	98.6		Mediur
5 P	ioneer 340	67.4			29.2	83	102.3		Mediu
7 P	roducers 315	66.9	66.3	1.1	28.4	81	99.9	102.0	Mediur
	rey 425	66.9			31.4	80	98.6		Mediur
	erris F-11	66.8			26.9	81	99.9		Mediu
0 P	P.A.G. b 366A	66.6	66.3	. 7	28.9	75	92.5	102.0	Mediur
1 F	unk G-29	66.4			30.6	89	109.7		Mediu
	llinois 1091A	66.2	65.2	1.8	31.2	80	98.6	100.6	Mediu
13 1	llinois 751	66.1	65.6	.9	30.4	79	97.4	101.3	Mediu
	DeKalb 458	65.9	65.6	. 4	28.2	77	94.9	101.3	Mediu
5 I	DeKalb 422	65.7	65.3	. 5	28.I	80	98.6	100.7	Mediu
	Ioosier Crost F-138	65.6	65.1	. 7	28.4	76	93.7	100.3	Mediu
	DeKalb 615	65.6	64.2	2.7	28.9	82	101.1	99.1	Mediui
	Vational 114-1	65.3			30.1		107.3		M-high
9 V	Vard 115A	64.8			29.2	77	94.9		Mediu
0° 1	llinois 101	63.9	63.7	. 3	27.9	86	106.0	98.4	Mediu
1d 1	llinois 269	63.3			30.1	80	98.6		Mediu
	roducers 320	63.1	62.9	. 3	28.9	81	99.9	97.0	Mediu
	owe 15	63.1	1111		29.3	82	101.1		Mediu
	PeKalb 404A	62.7	62.3	. 9	27.0	85	104.8	96.2	Mediu
	Crow 360	62.6	1111	• • •	33.5	69	85.1	11111	M-high
	Iorgan M-105	62.5	62.0	. 9	30.7	84	103.6	95.7	Mediu
	Moews 14	62.3			29.1	85	104.8		Mediu
28 C	Crow 514(W)	59.6			27.3	78	96.2		M-high
	Average of all entries	65.4	64.80	.9	29.1	81.1			
						~			

A difference of less than 3.8 bushels between total yields of any two entries in this summary is not significant.

		-		•					
			ock	Woodst	LTS:	RESUI	947 F	1	
5.1 Medium	125.1	114.1	93	22.8	. 4	51.9	52.3	Producers 305	1
1.0 Medium	121.0	90.8	74	24.5	. 7	50.2	50.7	Pioneer 349	2
9.5 Medium	119.5	103.1	84	26.3	.8	49.6	50.1	Furr 67A	3
5.4 Medium	115.4	106.7	87	25.1	.6	47.9	48.3	Moews 15	4
5.2 Medium	115.2	94.5	77	26.7	. 6	47.8	48.2	Hoosier Crost F-138	5
5.2 Medium	115.2	77.3	63	24.5	. 6	47.8	48.2	Huebsch H-44	5
4.4 M-low	114.4	100.6	82	23.9	1.2	47.5	48.1	DeKalb 410	7
4.2 Medium	114.2	103.1	84	23.6	. 4	47.4	47.6	Sieben S-450	8
1.3 Medium	111.3	100.6	82	25.7	2.2	46.2	47.3	Furr 66A	9
6.5 Medium	106.5	98.1	80	27.9	6.5	44.2	47.3	Crow 407	9
0.0 Medium	110.0	104.3	85	25.0	2.0	45.6	46.5	P.A.G. 60	11
0.6 Medium	110.6	98.1	80	24.8	. 6	45.9	46.2	Hunt 60(W)	12
7.5 M-low	107.5	81.0	66	27.0	3.2	44.6	46.2	Pioneer 343	12
Medium		90.8	74	27.2			46.0	Ward 115B	14
	109.2	104.3	85	27.3	1.2	45.3	45.9	Munson MX	15
4.6 Medium	104.6	114.1	93	30.0	4.5	43.4	45.6	Doubet D-1	16
7.0 Medium	107.0	98.1	80	25.9	2.4	44.4	45.4	P.A.G. 54	17
M-low		110.4	90	21.2			45.4	P.A.G. 299	17
Medium		100.6	82	26.3			45.2	Ferris F-11	19
2.4 Medium	102.4	103.1	84	29.1	5.9	42.5	45.1	DeKalb 615	20
7.7 Medium	107.7	94.5	77	26.3	. 7	44.7	45.0	DeKalb 458	21
5.7 Medium	106.7	89.5	73	25.4	1.3	44.3	44.9	Bear OK-20	22
5.5 Medium	106.5	89.5	73	27.6	1.4	44.2	44.8	Hoosier Crost FD-3	23
Medium		110.4	90				44.8		23
	107.0	104.3	85	25.7	. 4	44.4	44.7		25
7.0 Medium	107.0	100.6	82	23.9	. 4	44.4	44.5	DeKalb 422	26
		108.0	88	28.3			44.5	DeKalb 609	26
17	10	89.5 110.4 104.3 100.6	73 90 85 82	27.6 28.5 25.7 23.9	1.4	44.2 44.4 44.4	44.8 44.8 44.7 44.5	Hoosier Crost FD-3 Funk G-29 Illinois 101 (Producers) DeKalb 422	23 23 25 26

(Table is concluded on next page)

Table 4. - NORTHERN ILLINOIS - concluded

		Acre	-yield	Damaged corn in	ture in	Erect		g for—	– tive
Ran	k Entry			- shelled samples	grain at harvest	plants	Erect plants	Sound yielda	height of ear
	1947 R	ESUL	TS:	Woods	stock —	- concl	uded		
28 29 30	Illinois 1091A	43.7	bu. 42.3	perct. 4.0	perct. 30.5 34.9 27.5	perct. 75 50 86	perct. 92.0 61.3 105.5	perct. 101.9	Medium Medium Medium
31 32 33 34 35 35 35 38 39 40	Ward 115A. Lowe 52. Hoosier Crost F-140. Pioneer 340. Pioneer 344 Pioneer 4758 Lowe 32. Keystone 44. Pride D-66. Hulting 240.	43.2 43.1 43.0 42.7 42.7 42.7 42.7 42.5 42.4	42.8 42.5 42.2 41.4 	1.2 2.8 	29.8 25.9 27.2 28.5 28.8 30.0 27.2 30.1 30.5 29.5	76 83 86 78 85 90 85 92 88 84	93.2 101.8 105.5 95.7 104.3 110.4 104.3 112.9 108.0 103.1	103.1 102.4 101.7 99.8 	Medium Medium M-low Medium Medium Medium M-low Medium Medium
41 43 44 44 44 47 48 49	Producers 320 Producers 315 Producers 311 United U-36 Lowe 15 Frey 425 Producers 317 United U-41 Hulting 260 Funk G-16A	42.1 42.0 41.9 41.9 41.9 41.8 41.6 41.4	41.9 41.2 41.0 40.3 40.9	.4 2.4 2.6 4.1 1.6	29.3 28.4 25.3 26.5 28.8 31.7 27.9 31.0 27.6 28.5	83 79 84 90 92 78 81 92 75 94	101.8 96.9 103.1 110.4 112.9 95.7 99.4 112.9 92.0 115.3	101.0 99.3 98.8 97.1 98.5	Medium Medium Medium Medium Medium Medium M-high Medium Medium
51 52 52 54 54 54 54 58 58 59	Illinois 751 P.A.G. 270 P.A.G. 52 Producers 315 Nichols 5A DeKalb 404A Illinois 269 (Producers) Stiegelmeier S-9H Furr 44A Hoosier Crost FD-4	. 41.1 . 41.0 . 41.0 . 41.0 . 41.0 . 41.0	40.2 40.6 40.3 40.2 40.2 40.0 38.9 39.4	1.9 1.0 2.0 2.0 2.0 2.4 	30.1 30.1 23.5 33.1 30.0 26.3 29.8 33.7 30.1 30.0	74 84 88 79 92 90 77 95 69 87	90.8 103.1 108.0 96.9 112.9 110.4 94.5 116.6 84.7 106.7	97.0 96.4 97.1 97.0 95.9 93.7 94.9	Medium Medium Medium Medium Medium Medium Medium Medium M-low
61 63 64 65 66 66 68 69 70	Farmcraft PC-43. Morgan M-105 Huebsch H-55 Moews 85. Ward 110 Hoosier Crost F-145. Moews 14. Producers 510 National 115A Null N-32.	39.9 39.8 39.6 39.4 39.1 39.1 38.7 38.3	39.6 39.2 38.5 39.6 38.6 37.3 37.3 36.7 36.8	.4 1.8 3.5 .6 2.2 4.7 3.8 3.8 3.1	33.0 28.3 27.9 27.3 28.7 27.3 28.7 32.7 30.8 30.7	51 84 86 72 90 81 89 87 86 88	62.6 103.1 105.5 88.3 110.4 99.4 109.2 106.7 105.5 108.0	95.4 94.4 92.8 95.4 93.0 89.9 89.9 88.4 88.7	Medium Medium Medium M-low Medium Medium Medium Medium Medium
71 71 73 74 75 76 77 78 79	Furr 33. Pioneer 4040. Hoosier Crost F-150. Lowe 4(W). Ainsworth X-23. P.A.G. 366A Sieben S-340. Farmcraft FC-40. Lowe 34.	. 37.6 . 37.3 . 36.5 . 36.3 . 36.0 . 35.3	36.9 36.9 37.2 36.2 35.7 33.7 34.6	1.8 1.7 .3 .9 1.6 4.6 1.5	28.4 35.3 32.5 22.8 30.1 32.2 35.9 35.0 34.1	79 77 75 84 82 70 78 87 77	96.9 94.5 92.0 103.1 100.6 85.9 95.7 106.7 94.5	88.9 89.6 87.2 86.0 81.2 83.4	Medium Medium Medium Medium Medium Medium Medium M-low Medium
80 81	Crow 514(W)	. 33.3	32.7 41.5	1.6 2.0	27.3 25.7 28.4	76 73 81.6	93.2 89.5	78.8	Medium Medium

A difference of less than 5.0 bushels between total yields of any two entries in 1947 is not significant.

^{*} In 1947 accurate determination of the amounts of damage, sound yields, and sound yield scores could not be made for the entries which are blank. b Formerly Pfister. c In 1945 and 1946 seed was furnished by various growers; in 1947 yield was that of Illinois 101 (Producers). d In 1945 and 1946 seed was furnished by various growers; in 1947 yield was that of Illinois 269 (Producers).

Table 5. - CORN BORER DAMAGE: Mt. Morris, Kings, and Woodstock

Ranl	k Entry		ts broken ow ear*	Rank			Entry		nts broker low ear*
	SUMMARY:	Mt. Mor	ris 1943	3; Ki	ngs	1946;	Woodstock	1947	7
			perct.						perct.
1	Doubet D-1		2.8	11b	Produ	icers 32	0		6.8
	Moews 14		3.2	12	DeKa	alb 404	١		6.9
3	Funk G-29		3.4	13c	Illino	is 101			7.3
	Hoosier Crost F-140		3.9	14	DeKa	alb 422.			8.0
	Producers 510		4.3	15	DeKa	lb 615.			8.4
	Nichols 5A		5.0				t F-138		8.9
	DeKalb 458		5.7	17	Crow	514 (W	7)		9.3
	Furr 44A		5.7	18	Pione	er 340.			9.6
9	Illinois 751		5.9	19	Crow	360			9.8
10	Lowe 15		6.2		A	verage	of all entries		6.4

There are no significant differences between entries in this summary.

1947 DAMAGE

-		perct.		perci
1	Doubet D-1	. 6	42	Illinois 269 (Blackhawk) 7.9
2	United U-41	. 7	4.3	Sieben S-450 8.1
2	National 115A	. 7	44	DeKalb 410 8.2
4	Funk G-29	1.4	44	Null N-32 8.2
5	Pioneer 4758	2.0	46	Lowe 32 8.3
6	Lowe 34	2.4	47	Ward 110 8.5
6	Funk G-16A	2.4	47	Pioneer 343 8.5
8	Huebsch H-55	2.6	49	Furr 66A 8.6
8	Keystone 44	2.6	50	Huebsch H-44. 9.0
10	Moews 15	2.8		710
	2.200.00 2011111111111111111111111111111	2.0	50	Producers 311 9.0
11	P.A.G. 299	2.9	52	Bear OK-20 9.2
12	Nichols 5A	3.2	53	Ward 115B 9.4
12	Pioneer 4040	3.2	54	Moews 85 9.7
14	Lowe 15	3.5	55	DeKalb 458
15	United U-36	3.6	56	P.A.G. 54
16	Ward 115A	3.7	56	Hoosier Crost F-145 10.3
17	Producers 510	4.1	58	Farmcraft PC-43 10.6
18	Producers 305	4.3	59	Furr 44A
18	Producers 317	4.3	60	Producers 315
18	Hoosier Crost FD-4	4.3	00	Troducers 515
10	Hoosier Crost PD-4	4.5	61	Hoosier Crost F-138 11.6
21	Sieben S-340	4.9	62	Furr 67A
22	Pioneer 349	5.0	63	Crow 407
22	Stiegelmeier S-9H	5.0	64	Producers 320
22	Ainsworth X-23	5.0	65	P.A.G. 366A
22	Moews 14	5.0	66	Farmcraft FC-40
22	Illinois 1091A	5.0	67	DeKalb 422
27	National 114-1	5.4	68	Illinois 751
28	DeKalb 404A	6.0	68	P.A.G. 60
29	Morgan M-105	6.1	68	Pioneer 340
30	DeKalb 609	6.3	00	Fioneer 340 13.3
30	Dekaib 009	0.3	71	Hunt 60 (W)
30	Ferris F-11	6.3	72	DeKalb 615
32	Pride D-66	6.7	73	Crow 514 (W)
33		6.8	74	Illinois 101 (Blackhawk) 14.1
34	Hulting 240 Hoosier Crost F-140	6.9	75	Lowe 4 (W)
34	P.A.G. 270	6.9	76	Furr 33
34	Frey 425	6.9	77	Hoosier Crost FD-3
34		6.9	78	
38	Munson MX	7.0	78	Hulting 260
	Lowe 52			
39	P.A.G. 52	7.2 7.3	80	Lowe 6 (W)
40	Producers 315	7.6	81	Crow 360
41	Pioneer 344	7.0		Average of all entries 8.4
			1	

A difference of less than 8.0 in 1947 percentages is not significant.

a Includes only plants broken below the ear at point of damage by the borer, Pyrausta nubilalis (Hbn.).
b Prior to 1947 Producers 510 was Producers 1010 and Producers 320 was Producers 1020.
Average is for 1943 and 1947.
c 1947 seed of Illinois 101 is from Blackhawk Cooperative Hybrid Corn Association.

Table 6. — WEST NORTH-CENTRAL ILLINOIS: Galesburg

		Aore	-yield	Damaged corn in	Mois- ture in	Erect	Ratin	g for—	Compara – tive
Ran	k Entry		Sound	- shelled sample	grain at harvest	plants	Erect plants	Sound yield	height of ear
	S	UMM	ARY:	1945, 1	.946, an	d 1947			
		bu.	bu.	perct.	perct.	perct.	perct.	perct.	
1	Pioneer 339	. 87.1	86.4	1.0	21.4	62	104.0	105.8	M-high
	Pioneer 304		86.3	. 4	25.1	51	85.6	105,6	M-high
3	Doubet D-72	85.8	85.3	.6	22.3	63	105.7	104.4	M-high
4	Kelly K-374	85.1	83.9	1.7	21.4	56	94.0	102.7	M-high
5	Holmes Utility 39	, 84.7	84.0	.9	24.1	56	94.0	102.8	M-high
	P.A.G.* 5897		84.4	. 3	21.8	59	99.0	103.3	Mediun
	Crow 633		84.2	. 5	22.3	61	102.3	103.0	Mediun
	Holmes Utility 29		83.6	. 7	21.6	60	100.7	102.3	M-high
	DeKalb 847		82.5	1.6	21.4	68	114.1	101.0	M-high
10	Morton M-12	83.6	83.0	.9	22.9	65	109.1	101.6	M-high
	Funk G-169		82.9	.8	22.3	63	105.7	101.5	M-high
	Sieben S-440		82.9	.8	21.2	60	100,7	101.5	Mediun
	Farmcraft FC-47		82.5	. 7	21.6	50	83.9	101.0	Mediun
14	DeKalb 800A	. 82.8	80.2	3.2	23.3	58	97.3	98.2	Mediun
15	Null N-54	. 82.7	81.9	1.2	22.2	59	99.0	100.2	M-high
16 ^b	U. S. 13	. 82.5	81.6	1.1	23.3	65	109.1	99.9	M-high
17	Funk G-37	. 82.3	81.7	. 7	21.1	61	102.3	100.0	M-high
18	DeKalb 816	. 82.2	81.5	9	23.0	65	109.1	99.8	M-high
19 20	DeKalb 628A	82.0	80.8	1.6	22.8	62 68	104.0 114.1	98.9 98.6	M-high
	Moews 550	81.2	80.6 80.3		20.8 22.9	55	92.3	98.0	Mediun M-high
20	National 125	01.2	80.3	1.6	22.9	33	92.3	98.3	M-mgn
22	Crow 607	. 81.1	80.4	1.0	23.4	51	85.6	98.4	M-high
	Ward 120A		80.3	.9	23.0	51	85.6	98.3	M-high
24	Doubet D-42	. 80.0	78.7	1.9	22.9	63	105.7	96.3	M-high
25	Lowe 520	. 78.7	78.3	. 6	23.2	63	105.7	95.8	M-high
26	Keystone 42	. 77.5	77.1	. 4	21.9	57	95.6	94.4	M-high
27	Producers 311	77.1	76.5	.8	20.8	59	99.0	93.6	Mediun
28	Morgan M-546	76.9	75.8	1.4	23.0	58	97.3	92.8	M-high
	Average of all entries	82.5	81.7	1.0	22.4	59.6			
	and the contract of the contra	02.0	~			07.0			

A difference of less than 4.3 bushels between total yields of any two entries in this summary is not significant.

			1947	RESU	LTS				
1	Pioneer 313B	92.7	90.0	2.9	23.9	51	80.8	125.7	Medium
2	Schwenk S-24	87.3	85.9	1.5	22.2	62	98.3	120.0	Medium
3	Pioneer 336	86.0	85.1	1.1	21.9	65	103.0	118.8	M-high
4	Stiegelmeier S-201	84.4	80.5	4.5	20.9	66	104.6	112.4	Medium
5	Ward 120A	83.2	81.3	2.0	22.9	43	68.1	113.5	Medium
6	Pioneer 339	82.8	81.6	1.2	20.9	61	96.7	114.0	Medium
7	Stiegelmeier S-301	81.8	78.3	4.3	21.1	87	137.9	109.4	Medium
8	P.A.G. 390	80.9	79.3	2.2	20.1	65	103.0	110.7	Medium
9	Bear OK-50	81.0	80.3	.6	22.2	67	106.1	112.2	Medium
9	Pioneer 304	81.0	80.2	1.0	27.7	47	74.5	112.0	Medium
11	Producers 730	80.7	78.7	2.5	23.6	43	68.1	109.9	Medium
12	Holmes Utility 39	80.3	79.0	1.6	26.0	55	87.2	110.3	Medium
13	Morgan M-600	79.7	77.6	2.8	22.9	81	128.4	108.4	Medium
14	Pioneer 300	79.5	78.1	1.8	23.5	72	114.1	109.1	Medium
14	U. S. 13 (Morton)	79.5	76.8	3.5	23.4	73	115.7	107.3	Medium
14	DeKalb 800A	79.5	74.9	5.9	22.4	49	77.6	104.6	Medium
17	U. S. 13 (Lepper)	79.0	77.8	1.6	24.4	63	99.8	108.7	Medium
17	Kelly K-374	79.0	76.6	3.0	21.9	61	96.7	107.0	Medium
19	Producers 709	78.9	77.8	1.3	23.3	52	82.4	108.7	Medium
20	Lowe 510	78.5	77.6	1.2	21.5	64	101.4	108.4	Medium
21	Doubet D-72	78.2	77.3	1.4	22.1	75	118.8	108.0	Medium
22	Crow 633	77.7	77.0	1.0	23.8	56	88.7	107.5	Medium
23	Kelly K-100	77.6	76.9	.8	21.8	54	85.6	107.4	M-high
24	Hulting 380	77.3	75.5	2.4	22.2	63	99.8	105.4	Medium
25	Ainsworth X-21	76.8	75.8	1.4	23.3	85	134.7	105.9	Medium
25	DeKalb 847	76.8	73.8	3.9	22.6	82	129.9	103,1	Medium
27	Bear OK-40	76.0	74.9	1.5	20.7	60	95.1	104.6	Medium

(Table is concluded on next page)

Table 6. — WEST NORTH-CENTRAL ILLINOIS — concluded

n .	-	Acre	-yield	Damaged corn in	Mois- ture in	Erect	Ratin	g for—	Compara-
Rank	Entry		Sound	— shelled	grain at harvest	plants	Erect plants	Sound yield	height of ear
		1947 I	RESU	JLTS —	conclu	ded			
28 A	Bear OK-60hinsworth X-23	bu. 75.9 75.9 75.9	bu. 75.1 75.0 74.4	perct. 1.0 1.2 1.8	perct. 23.8 21.2 22.1	perct. 52 67 66	perct. 82.4 106.1 104.6	perct. 104.9 104.7 103.9	Medium Medium Medium
31 S 33 I 34 I 35 F 36 N 37 N 37 I 39 F	owe 24 jieben S-440 llinois 972-1 (Pringle) llinois 21 (Burrus) Producers 940 vational 125-1 vall N-54 beKalb 817A Holmes Utility 29 Hulting 366	75.7 75.7 75.4 75.3 75.2 75.0 74.5 74.5 74.3	74.8 74.4 74.1 74.0 73.4 71.8 72.3 71.7 72.9 73.2	1.2 1.7 1.7 1.8 2.2 4.3 3.0 3.7 1.7	21.3 20.8 26.2 21.3 21.2 21.9 22.2 22.0 22.5 22.6	66 69 36 66 83 57 71 60 55	104.6 109.4 57.0 104.6 131.5 90.3 112.5 95.1 87.2 110.9	104.5 103.9 103.5 103.4 102.5 100.3 101.0 100.1 101.8 102.2	Medium Medium M-high Medium Medium Medium Medium Medium Medium
42 II 43 M 44 M 45 II 46 II 47 F 48 M 49 M	Jnited U-65. Joubet D-42. Joews 520. Joews 520. Joews 514. Joek 318 16. Joek 318 16. Joek 3897. Joek 380. Joek 3	73.4 73.1 72.7 72.4 72.3 72.1 72.0 71.9 71.8 71.8	71.0 72.0 72.0 69.8 71.0 70.8 71.5 71.3 70.7	3.4 1.5 .9 3.7 1.7 1.8 .8 1.7 2.0	21.9 21.4 22.2 20.8 21.0 21.7 22.0 22.2 22.9 22.6	54 74 84 60 53 70 68 80 71 56	85.6 117.3 133.1 95.1 84.0 110.9 107.8 126.8 112.5 88.7	99.2 100.6 100.6 97.5 99.2 98.9 99.9 99.6 98.7 98.3	Medium Medium Medium Medium Medium Medium Medium Medium Medium
52 M 53 F 54 L 54 H 56 M 57 F 57 S 59 H	DeKalb 628A. Munson M-19 Producers 315 Lowe 523 Hulting 101 Morgan M-546 Purr 67 Stewart S-8 Funk G-37	71.3 70.8 70.7 70.6 70.6 70.3 70.1 70.1 69.3 69.1	69.2 67.5 70.2 70.0 69.5 67.8 69.2 68.7 65.5 67.9	2.8 5.1 .6 1.2 1.8 3.4 1.3 2.0 5.5	21.8 22.0 20.3 24.4 21.1 24.9 20.7 24.8 22.0 21.4	64 65 74 43 54 53 48 41 40 58	101.4 103.0 117.3 68.1 85.6 84.0 76.1 65.0 63.4 91.9	96.6 94.3 98.0 97.8 97.1 94.7 96.6 95.9 91.5 94.8	M-high Medium Medium Medium M-high M-low M-high Medium Medium
62 C 63 U 64 F 65 F 66 V 66 H 68 F 69 M	Moews 550. Frow 607. J. S. 13 (Morgan) A.G. 170. Farmcraft FC-47. Vard 115B. Hulting 260. Ferris F-A1. Moews 14. United U-59.	68.5 68.1 67.1 66.7 66.4 66.3 66.3 66.2 65.9	67.2 66.6 65.1 66.1 65.3 65.4 65.2 66.1 65.6	1.6 2.3 3.0 .9 1.8 1.2 1.6 .1	20.3 24.3 24.4 20.8 22.0 20.6 22.2 21.6 20.4 27.2	82 45 64 80 41 55 82 56 45 64	130.0 71.3 101.4 126.8 65.0 87.2 130.0 88.7 71.3 101.4	93.8 93.0 90.9 90.9 91.2 91.3 91.1 92.3 91.6 90.8	Medium Medium Medium Medium Medium Medium Medium Medium Medium
72 F 73 F 73 F 75 F 76 F 77 N 78 F 79 K 80 F	P.A.G. 392. Producers 311 Hoosier Crost 746. P.A.G. 299. Huey H-42. Huey H-20. Morton M-33. Hoosier Crost F-169. Eveystone 42. Huey H-50. Lowe 520. Average of all entries.	64.8 64.1 63.0 63.0 62.8 61.1 61.0 58.5 58.3 56.8 54.6	64.0 63.1 62.3 62.2 60.8 59.8 59.6 56.5 57.9 56.0 54.1 71.6	1.1 1.6 1.3 1.5 3.4 2.1 2.6 3.5 .6 1.2 1.2	20.7 20.4 23.2 20.4 20.4 22.8 22.8 22.6 21.5 22.3 22.9 22.3	70 58 69 59 80 74 76 63 60 84 75	110.9 91.9 109.4 93.5 126.8 117.3 120.4 99.8 95.1 133.1 118.9	89.4 88.1 87.0 86.9 84.9 83.5 83.2 78.9 80.9 78.2 75.6	Medium Medium Medium Medium Meligh Medium Medium Medium Medium Medium Medium Medium

A difference of less than 6.0 bushels between total yields of any two entries in 1947 is not significant.

 $^{^{\}rm a}$ Formerly Pfister. $^{\rm b}$ 1947 yield computed as average of U. S. 13 seed from the following producers: Lepper, Morgan, and Morton.

Table 7.—HYBRID RESISTANCE TO CORN ROOTWORM*

DAMAGE: Galesburg Summary, 1943 and 1947

Rank	Entry	Plants leaning 30 degrees or more	Plants leaning more than 45 degrees	Resistance rating com- pared with average
		perct.	perct.	
- 1	Lowe 520	6.7	1.7	243
2	DeKalb 816		1.6	203
3	DeKalb 800A		2.0	200
4	Morgan M-546		2.0	197
5	Doubet D-72	9.2	1.8	194
6	Moews 550	10.0	2.5	165
7b	Producers 940	13.8	1.4	149
8	DeKalb 817A	12.8	2.6	138
9	Funk G-37	10.5	5.3	136
10	Illinois 21	13.9	4.2	111
11	DeKalb 628A	15.6	3.9	106
12	Pioneer 339	18.4	2.6	105
13	P.A.G. 6 5897	14.1	4.8	104
14	Crow 633	16.5	3.7	103
15^{d}	National 125		4.1	102
16	U. S. 13	16.1	6.0	88
17	Farmeraft FC-47	22.8	4.7	77
18	Hoosier Crost 668	18.7	8.4	70
19°	Illinois 972	24.2	8.6	60
20	Crow 607	26.5	7.6	59
20	Moews 14	25.1	8.4	59
22b	Producers 730	25.8	11.6	41
	Average of all entries		4.6	100

Differences between hybrids are not significant.

Mostly southern corn rootworm, Diabrotica duodecimpunctata (F.).
 b In 1943 entries were Producers 1040 and 1030 respectively.
 e Formerly Pfister.
 d In 1947 entry was National 125-1.
 e In 1947 entry was Illinois 972-1 (Pringle).

Table 8.— HYBRID RESISTANCE TO CORN ROOTWORM DAMAGE: Galesburg, West North-Central Illinois, 1947

Rank	nk Entry	Plants leaning 30 degrees or more	Plants leaning more than 45 degrees	Resistance rating com- pared with average ^b	Rank	Entry	Plants leaning 30 degrees or more	Plants leaning more than 45 degrees	Resistance rating com- pared with average ^b
		perct.	perct.				perct.	perct.	
-	Lowe 520	2.2	0	2700	43	Hoosier Crost 746	29.7	14.3	102
2	Huev H-42	8.9	0	874	43	Lowe 514.	30.3	14.1	102
3	Huev H-50	9.5	0	632	45	Hulting 366.	35.3	11.8	101
4		13.5	0	443	45	Pioneer 300	42.2	8.3	101
S	Stiegelmeier S-301.	6.7	3.2	371	47	Munson M-19.	31.6	14.3	66
9	Morton M-12	17.0	0	349	47	DeKalb 628A.	33.0	13.6	66
7	Ainsworth X-21	. 11.8	2.7	345	49	Holmes Utility 29	36.2	12.4	46
∞	Lowe 510.	18.2	0	326	49	Farmcraft FC-47	44.2	8.4	26
6	Producers 315.	10.2	4.6	306	49	Hulting 101	37.6	11.9	97
10	P.A.G. 170	18.4	1.1	288	52	Munson M-15	36.6	12.9	95
2	Huev H-20	20.7		288	25	DeKalb 817A	40.9	10.0	95
13	Morgan M-600	21.7	· C	275	4	Pioneer 330	43.1	101	04
7 2	Doubet D-72	0 81	-	262		Huay H.73	38.5	13.1	00
2.	Doubet D-12.	0.00		261		11 C 12 (Monton)	21.5	1. 71	7.0
* :	Doubet D-42	0.07	0.0	107		O. S. 13 (Morton)	01.0	7.01	76
5	Moews 520.	0.61	3.0	238		Schwenk S-24	53.0	5.5	82
10	Funk G-169	0.91	5.7	217		Illinois 21 (Burrus)	37.4	17.2	83
17	Null N-54	. 19.4	4.3	212		Hulting 380	2.99	5.9	82
18	P.A.G. 392	. 22.0	3.3	208	9	Crow 633	40.2	16.5	81
19	DeKalb 816	. 15.5	8.9	205	61	U. S. 13 (Lepper)	35.3	19.6	80
61	Sieben S-440.	. 18.8	5.1	205	62	National 125-1	41.2	17.6	28
21	Bear OK-40	23.1	3.8	194	63	DeKalb 847	41.7	17.6	7.7
21	Hulting 260	30.7	0	194	49	Funk G-37	35.5	22.4	74
2.3	Moews 550	15.7	×	179	5	P A G 5807	47.9	10.1	69
24	Producers 940	25.8	4	175	9	Crow 607	46.5	20.8	67
25	Stiegelmeier S. 201	21.8	4	172	99	P A G 300	8 1 1 8	23.6	67
26	Producers 709	27.8	4	165	8	Ward 120A	46.4	22.7	55
27	DeKalb 800A	9 01	× ×	163	9	Pioneer 313R	50.5	20.5	45
28	Lowe 24	× × ×	ı,	162	9	Producers 311	20.02	4 %	4.
20	Rear OK-50	31.2		150	7.	United IL-65	2 92	27.0	5.3
30	Pioneer 304	20.2	200	22	22	Hoosier Crost F.160	× 4.5	20.8	52
31	Kelly K-100	28.3	4	152	12	Holmes Phility 30	45.8	34 6	5.5
32	United IL-50	21.2		121	14	Ward 115B	70.7	27.3	47
33	Pioneer 336	23.5	20,	145	7.4	Moews 14	27.6	34.	47
34	Keystone 42	26.0	1.7	143	76	Harris H.A.1	62.3	34 0	45
3.5	Morgan M 546	2,02		130	7.0	Illinois 079 1 (Deingle)	5.79	22.2	24
35	Moston M 200	† · · · · ·	7.0	139	9 6	IIIIIIIIII 9/2-1 (FIIIIBIE)	7.7	33.3	7.7
30	A immediate V 22	7.07	0.0	130	0 5	Hoosier Crost boo	07.0	30.0	7.5
300	AIIISWOTHII A-63	7.07	ć.	001	2 6	Stewart 3-0	1.00	7.00	77.
20	Lowe 525	200.2	0.0	071	8	Bear OK-00.	8.71	6.04	27
66	D A C 200	0.67	, ;	671	01	Froducers /30	6.47	7:44	10
7 =	Fig. 6.299	22.7	11.5	119		Average of all entries	33.6	12.9	100
47	Morton M-33	30.0	12.5	102					
1	MOLION ALTONOS SERVICES SERVIC	0.00	5.0	cot					

In Column 1 a difference of less than 33.6 per cent between any two entries is not significant.

* Especially southern corn rootworm, Diabrotica duodecimpunctata (F.). b High rating indicates better standing ability.

Table 9. — EAST NORTH-CENTRAL ILLINOIS: Sheldon

-		Acre	-yield	Damaged corn in	Mois- ture in	Erect	Ratin	g for—	Compara-
Ran	k Entry		Sound	shelled sample	grain at harvest	plants	Erect plants	Sound yield	height of ear
	SI	JMM.	ARY:	1945, 1	.946, an	d 1947			
		bu.	bu.	perct.	perct.	perct.	perct.	perct.	
1	Pioneer 313B	91.7	90.7	1.2	25.8	69	95.8	109.0	Medium
2	Pioneer 304	88.0	87.1	1.0	26.8	68	94.4	104.7	Medium
3	Morton M-380	87.9	87.3	. 7	24.9	71	98.6	104.9	Medium
3	Frey 644	87.9	87.3	.8	25.0	71	98.6	104.9	M-high
5 6	Pioneer 332	87.6 86.5	87.2 85.3	.5 1.6	26.5 25.0	70 74	97.2 102.8	104.8 102.5	M-high M-high
7	Frey 692 Pioneer 300	86.0	85.5	.5	24.3	74	102.8	102.3	M-high
8	Keystone 38	85.7	85.4	. 4	24.3	76	105.6	102.6	M-high
9	DeKalb 847	85.6	84.7	1.2	23.6	73	101.4	101.8	Medium
ý	Producers 730	85.6	84.7	1.2	25.1	72	100.1	101.8	M-high
1	Doubet D-47	85.4	84.8	. 7	24.3	76	105.6	101.9	M-high
2	DeKalb 800A	85.3	84.4	1.2	24.7	75	104.2	101.4	Mediun
3	Crow 607	84.6	83.5	1.4	24.6	69	95.8	100.4	M-high
4	Kelly K-374	84.3	83.7	. 8	23.2	71	98.6	100.6	M-high
5	Funk G-94	84.0	83.5	.6	25.5	75	104.2	100.4	M-high
6ª	U. S. 13	83.9	82.8	1.4	25.0	71	98.6	99.5	M-high
6 8	Kelly K-77 Producers 940	83.9 83.8	82.4 82.8	$\frac{1.7}{1.3}$	23.9 23.1	73 77	101.4 106.9	99.0 99.5	M-high Mediun
9	Lowe 520	83.7	83.5	.2	27.1	65	90.3	100.4	M-high
9	Stiegelmeier S-360	83.7	83.2	.5	22.2	50	69.4	100.0	Medium
1	Morton M-33	83.6	82.9	.9	24.7	76	105.6	99.6	M-high
2	DeKalb 628A	83.3	83.0	. 4	24.7	72	100.1	99.8	M-high
3	Frey 645	81.9	81.3	. 8	24.1	76	105.6	97.7	Mediun
4	Farmcraft FC-69	81.8	81.5	. 4	25.8	71	98.6	98.0	M-high
5	Crow 607(W)	81.3	81.0	. 4	23.7	68	94.4	97.4	M-high
6	Crow 608	80.8	80.1	.9	24.0	75	104.2	96.3	M-high
7 8	Hoosier Crost 668	80.4	80.1	.4	23.8	70	97.2	96.3	M-high
8 9	DeKalb 817A	80.0	78.9 78.5	1.5 1.5	24.8 22.5	72 80	$100.1 \\ 111.1$	94.8 94.4	Mediun Mediun
0	National 118 Crow 633	79.6 77.5	77.3	.3	25.4	74	102.8	94.4	Mediun
1	Hoosier Crost F-170	77.2	76.3	1.1	25.4	75	104.2	91.7	Mediun
	Average of all entries	84.0	83.2	.9	24.6	71.9			

A difference of less than 4.2 bushels between total yields of any two entries in this summary is not significant.

			1947	RESU	LTS				
1	Pioneer 304	82.0	81.5	.6	25.8	81	97.8	116.1	Medium
2	Bear OK-50	80.7	78.0	3.2	23.0	83	100.2	111.1	Medium
3	Pioneer 313B	79.6	77.8	2.2	27.4	83	100.2	110.8	Medium
4	U. S. 13 (Morton)	79.1	79.0	. 1	25.9	83	100.2	112.7	Medium
5	DeKalb 847	78.5	76.6	2.6	25.8	86	103.9	109.1	Medium
6	Frey 644	78.3	77.8	. 6	23.1	84	101.4	110.8	Medium
7	Pioneer 336	77.9	76.6	1.5	21.7	83	100.2	109.1	Medium
8	Schwenk S-66	77.2	76.3	.9	23.7	81	97.8	108.7	Medium
8	Kelly K-77	77.2	75.2	2.5	23.2	82	99.0	107.1	Medium
10	Huey H-50	77.0	76.8	. 4	24.7	86	103.9	109.4	Medium
11	Morton M-12	76.5	75.3	1.4	23.9	80	96.6	107.3	Medium
12	Pioneer 332	76.4	75.9	. 8	26.6	80	96.6	108.1	M-high
13	Producers 940	76.3	75.4	1.2	23.6	81	97.8	107.4	Medium
14	Illinois 21 (Mountjoy)	76.2	75.9	. 2	25.0	73	88.2	108.1	Medium
15	Farmcraft PC-63	76.1	75.1	1.3	22.4	77	93.0	107.0	M-low
15	Appl A-130	76.1	74.0	2.6	24.4	79	95.4	105.4	Medium
17	Ferris F-14	75.2	73.8	1.6	22.9	85	102.7	105.1	Medium
18	Daily DX9	75.0	74.9	. 1	24.0	83	100.2	106.5	Medium
18	Ward 115C	75.0	74.8	. 2	22.5	85	102.7	106.3	M-low
18	Keystone 38	75.0	74.6	.6	22.9	88	106.3	106.3	Medium
18	Stiegelmeier S-360	75.0	74.4	. 7	23.8	77	93.0	106.0	Medium
22	Ainsworth X-13-3	74.9	74.8	. i	26.1	84	101.4	106.5	Medium
22	Bear OK-88T	74.9	73.7	1.5	26.0	83	100.2	105.0	Medium
24	Bear OK-40	74.5	73.5	1.4	25.1	84	101.4	104.7	Medium

(Table is concluded on next page)

Table 9. — EAST NORTH-CENTRAL ILLINOIS — concluded

D	. E	Acr	e-yield	Damaged corn in	ture in	Erect		g for—	tive
Ran	k Entry	Tota	1 Sound	– shelled sample	grain at harvest	plants	Erect plants	Sound yield	height of ear
		1947	RESU	LTS—	conclu	ded			
		bu.	bu.	perct.	perct.	perct.	perct.	perct.	
24	Producers 730	74.5	73.0	2.0	25.0	83	100.2	104.0	Medium
	DeKalb 800A	74.4	72.9	2.2	23.8	81	97.8	103.8	Medium
27	U. S. 13 (Kelly)	73.5	73.3	. 3	23.1	76	91.8	104.7	Medium
27 29	Pioneer 300	73.5 73.3	$\frac{73.2}{72.0}$.4 1.3	23.5 26.6	88 83	106.3 100.2	104.3 102.6	Medium Medium
	Frey 645	73.3	70.8	3.3	27.2	88	106.3	100.8	Medium
1	Moews 523	73.1	72.4	.9	26.1	80	96.6	103.1	Mediun
31	Kelly K-374	73.1	71.6	2.0	23.9	77	93.0	102.0	Mediun
33	Kelly K-88	72.5	71.9	.9	23.8	83	100.2	102.4	Mediun
	Stiegelmeier S-380	72.4	72.3	2.1	24.1	84	101.4	103.0	Medium
	Illinois 1425	$72.4 \\ 72.0$	70.5 71.2	2.8	25.8 22.3	84 84	101.4	100.4 101.4	Medium
36 36	Sibley 777	72.0	69.0	$\frac{1.1}{4.1}$	25.7	86	101.4 103.9	98.3	Mediun Mediun
88	Producers 945	71.9	68.8	4.5	24.1	73	88.2	98.0	Mediun
	Ward 120A	71.6	71.2	.7	25.2	87	105.1	101.4	Mediun
	Ainsworth X-14A	71.6	69.9	2.5	30.8	84	101.4	99.6	M-high
39	Crow 607	71.6	69.5	3.1	25.0	84	101.4	99.0	Mediun
12	Morton M-380	71.5	71.1	.5	26.1	86	103.9	101.3	Mediun
43	Moews 550	$71.4 \\ 71.0$	70.5 70.9	1.1	21.6 26.7	76 90	$91.8 \\ 108.7$	$100.4 \\ 101.0$	Mediun
14 14	Illinois 972-1 (Appl) Seeber 11A	71.0	67.7	4.6	28.5	81	97.8	96.4	Mediun Mediun
16	P.A.G. 392	70.6	69.3	1.8	23.0	79	95.4	98.7	Mediun
17	Appl A-136	70.3	70.2	. 2	23.0	88	106.3	100.0	M-low
48	Appl A-136 Doubet D-47	70.2	69.8	. 6	25.9	82	99.0	99.4	Mediun
49	Powers 149	70.1	69.9	. 1	22.7	80	96.6	99.6	M-low
49	Crow 607(W)	70.1	69.6	. 5	25.3	86	103.9	99.1	Mediun
19	National 118	70.1	68.3	2.5	23.8	87	105.1	97.3	Mediun
52	Pioneer 339	69.8	69.5	.6	22.8	83	100.2	99.0	M-low
	Hoosier Crost 12	69.6	69.4	. 1	21.6	89	107.5	98.9 95.3	M-low
54 55	Lowe 555	69.2 69.0	66.9 68.6	3.4	$\frac{24.8}{22.0}$	72 88	87.0 106.3	95.3	M-low Mediun
55	P.A.G. 170 Hoosier Crost F-169	69.0	67.3	2.6	24.3	81	97.8	95.9	M-low
57	Hoosier Crost F-150	68.5	68.4	. 2	22.5	79	95.4	97.4	M-low
	P.A.G. 164	68.4	65.5	4.2	23.2	86	103.9	93.3	Mediun
59	P.A.G. 173	68.1	67.9	.3	29.7	86	103.9	96.7	Mediun
50	Huey H-42	67.9	67.1	1.0	25.1	87	105.1	95.6	Mediun
51	Moews 520	67.4	67.0	.6	24.5	83	100.2	95.4	Mediun
62	Hoosier Crost F-170	67.2	66.4	1.4	26.5	77	93.0	94.6	Mediun
53	Farmcraft FC-69	66.6	66.3	. 6	26.2	80	96.6	94.4	Mediun
64	Funk G-94	66.5	66.3	.6	25.6	84 84	101.4	94.4	Mediun
55 56	Trisler T-22 Lowe 514	66.4 66.1	65.8 65.8	1.0	25.3 26.0	86	$101.4 \\ 103.9$	93.7 93.7	Mediun Mediun
56	Funk G-211	66.1	65.2	1.2	23.5	87	105.1	92.9	Mediur
58	Funk G-211 DeKalb 628A	66.0	65.8	. 3	26.4	85	102.7	93.7	Mediun
59	Hoosier Crost F-145	65.4	65.0	. 5	21.9	82	99.0	92.6	M-low
70	DeKalb 817A	65.2	63.7	2.2	25.6	86	103.9	90.7	Mediun
71	Producers 311	64.8	64.5	.5	22.4	86	103.9	92.2	M-low
72 73	Illinois 1428 Lowe 520	64.6	64.4	. 2	$24.4 \\ 24.7$	84 83	101.4 100.2	91.7 91.7	Mediun Mediun
73 74	Crow 633	64.0	63.8	.1	24.7	77	93.0	90.9	Mediun
75	Hoosier Crost 668	63.3	62.8	,6	23.7	79	95.4	89.5	Mediun
76	Lowe 523	62.5	62.2	. 7	26.0	77	93.0	88.6	Mediun
77	Lowe 580	62.2	59.6	4.2	24.1	79	95.4	84.9	Mediun
78	Lowe 580	61.9	60.6	2.0	26.3	88	106.3	86.3	Mediun
79	Appl A-202	61.2	61.1	. 2	23.0	83	100.2	87.2	M-low
80	Embro 95	60.2	60.0	. 2	22.8	85	102.7	85.5	M-low
81 *	Crow 608	60.0	59.6	.6	25.5	86	103.9	84.9	Mediun
	Average of all entries	71.1	70.2	1.3	24.6	82.8			

A difference of less than 7.3 bushels between total yields of any two entries in 1947 is not significant.

^{* 1947} yield is computed as average of U. S. 13 from the following producers: Kelly, Morton, and Pfeifer.

Table 10. - CORN BORER DAMAGE: Milford and Sheldon

Rank		Entry	Plants broken below ears	Rank		ints broker elow eara
		SUMMARY:	Milford 19	944; S	Sheldon 1945 and 1947	
			perct.			perct.
1	Produce	ers 940	14.6	14	DeKalb 628A	19.7
2		20		15	Hoosier Crost 668	20.3
3	Crow 60	07	15.4	16	DeKalb 847	20.4
4 b		9722		16	Pioneer 332	20.4
5	Frey 64	4	17.5	18	Producers 730	21.0
6		M-380		19	DeKalb 800A	21.4
7		300		20	Pioneer 304	21.7
8		08		20	D: 226	24 7
9	Crow of	07 _. (W)	18.6	20	Pioneer 336	21.7
10	Stiegeln	neier S-360	18.8	22	Kelly K-374	22.6 23.0
10	11th air	21	18.8	23	Frey 645 Crow 633	23.8
10		D-47		25	DeKalb 817A	24.3
13		D-47		26	Funk G-94	25.5
10	0. 5. 10	·····	10.7	20	Average of all entries	19.7
		Differences be	etween en	tries	are not significant.	
			1947 D	AMA	GE	
1	Hoosier	Crost 12	7	41	Ainsworth X-13-3	4.5
2				41	Huey H-50	
3	Morton	202 M-33	9	41	Pioneer 304	. 4.5
4	P.A.G.	173	1.5	41	Farmcraft PC-63	4.5

		154/ 1	AMAGE	
1 2 3 4 4 6 6 6 6 9	Hoosier Crost 12 Appl A-202 Morton M-33 P.A.G. 173 P.A.G. 392 Funk G-211 Morton M-380 Farmcraft FC-69 Illinois 972-1 (Appl) Seeber 11A	.7 .8 .9 1.5 1.5 1.6 1.6 2.2 2.4	41 Ainsworth X-13-3 4.5 41 Huey H-50 4.5 41 Pioneer 304 4.5 41 Farmcraft PC-63 4.5 45 Bear OK-88T 4.7 45 Ward 115C 4.7 47 Ward 120A 4.8 48 U. S. 13 (Morton) 4.9 49 Morton M-12 5.0 50 Doubet D-47 5.1	
11 12 13 13 13 16 17 18 18	Appl A-130 Pioneer 332 Producers 311 Illinois 1425 Crow 608 Appl A-136 Embro 95 Hoosier Crost F-170 P.A.G. 164 Keystone 38	2.7 2.8 2.9 2.9 3.0 3.1 3.2 3.2 3.3	50 Frey 645 5.1 52 P.A.C. 170 5.2 52 Lowe 520 5.2 52 Hoosier Crost F-145 5.2 55 Funk G-94 5.3 56 Crow 607 5.6 56 Crow 607 (W) 5.6 58 DeKalb 800A 5.7 58 Lowe 555 5.7 60 Stiegelmeier S-360 5.9	
21 21 21 25 26 27 27 29	Kelly K-77 Pioneer 300. U. S. 13 (Pfeifer) Lowe 523. Frey 692 Ferris F-14 Producers 940. Kelly K-88 Bear OK-40 Moews 550.	3.4 3.4 3.4 3.5 3.6 3.7 3.8 3.8	61 Producers 730. 6.0 62 National 118. 6.1 62 Trisler T-22. 6.1 62 Hoosier Crost F-169. 6.1 65 DeKalb 817A. 6.3 66 Stiegelmeier S-380. 6.5 67 Illinois 21 (Mountjoy). 6.6 68 Moews 520. 6.7 69 Sibley 777. 6.8 70 Schwenk S-66. 7.3	
31 32 33 33 36 36 36 36 40	Frey 644 Pioneer 339 Pioneer 336 Bear OK-50 Lowe 514 Daily DX9 Moews 523 Hoosier Crost 668 DeKalb 847 DeKalb 628A	4.0 4.1 4.2 4.2 4.3 4.3 4.3 4.3	71 Producers 945. 7. 4 72 Ainsworth X-14A. 7. 5 73 Lowe 580. 7. 9 74 U. S. 13 (Kelly). 8. 0 75 Crow 633. 8. 1 76 Hoosier Crost F-150 8. 2 77 Kelly K-374. 8. 3 78 Illinois 1428. 8. 5 79 Powers 149. 8. 7 80 Huey H-42. 8. 9 81 Pioneer 313B. 10. 9 Average of all entries. 4. 8	

Differences between entries are not significant.

^{*} Includes only those plants broken below the ear at point of damage by the borer, *Pyrausta nubilalis* (Hbn.). b Entry included as Illinois 972-1 (Appl) in 1947, as Illinois 972-2 (Appl) in 1945, and Illinois 972-1 in 1944.

Table 11. — SOUTH-CENTRAL ILLINOIS: Sullivan

		Acre	-yield	Damaged corn in	Mois- ture in	Erect	Ratin	g for—	Compara tive
Rank	Entry		Sound	– shelled	grain at harvest	plants	Erect plants	Sound yield	height of ear
	S	UMM.	ARY:	1945, 1	946, an	d 1947			
		bu.	bu.	perct.	perci.	perct.	percl.	perct.	
1 I	Doubet D-41	90.1	89.5	. 6	21.2	83	91.8	110.1	M-high
	Bear OK-40 Producers 1050	89.4 88.9	89.0 88.5	.5 .6	19.8 21.2	92 87	101.8 96.2	109.5 108.9	Medium
	Illinois 21	88.6	87.2	1.7	20.5	93	102.9	107.3	M-high Medium
5 1	Pioneer 313B	87.8	86.7	1.2	21.9	93 87	96.2	106.6	Mediun
6 6	Crow 607	86.8	86.1	1.5	21.8	83	91.8	105.9	M-high
7 Ì	Pioneer 332	86.3	85.6	.9	23.5	92	101.8	105.3	M-high
8 I	Kelly K-374	86.0	83.9	2.5	19.7	89	98.4	103.2	Medium
	Morton M-12	85.9	85.2	. 8	21.0	91	100.7	104.8	Medium
10 I	Pioneer 300	85.8	83.9	2.4	21.1	94	104.0	103.2	Medium
	P.A.G.b 164	85.6	84.1	1.9	20.3	95	105.1	103.4	Mediun
12° I	Illinois 201	85.4	84.9	. 7	19.9	90	99.6	104.4	Mediun
	Producers 940	84.8	82.1	3.6	20.5	95	105.1	101.0	Mediun
	Farmcraft FC-81 Pioneer 336	$84.7 \\ 84.3$	83.6 83.5	1.3	20.7 20.7	91 94	100.7 104.0	102.8 102.7	Medium Medium
	J. S. 13	83.9	82.9	1.2	20.7	91	100.7	102.7	Mediun
17 Î	DeKalb 816	82.6	81.2	1.6	20.7	91	100.7	99.9	Medium
18 Î	Keystone 38	82.4	81.6	1.0	21.2	92	101.8	100.4	Medium
19 (Crow 805	81.0	80.7	. 5	21.4	93	102.9	99.3	Mediun
19 V	Ward 120A	81.0	80.4	.8	21.5	92	101.8	98.9	M-high
	Crow 608	80.9	80.1	1.2	20.6	94	104.0	98.5	Medium
	National 125-1	80.9	79.0	2.7	20.4	93	102.9	97.2	Medium
23 F	Kelly K-99	80.5	79.5	1.3	20.6	88	97.3	97.8	Mediun
24f 1 25 F	llinois 126	80.0 79.6	79.7	.4	20.8	89 94	98.4 104.0	98.0 97.3	Medium
	Hoosier Crost 746 Morgan M-546	79.0	79.1 78.1	.5 1.3	21.1 21.4	91	104.0	96.1	Mediun M-high
	P.A.G. 612(W)	78.8	77.9	1.5	23.1	84	92.9	95.8	M-high
	Ilinois 972A-1	77.8	75.6	2.7	21.6	93	102.9	93.0	Medium
	Pioneer 505(W)	76.8	75.4	2.1	23.4	89	98.4	92.7	M-high
30 V	Whisnand 917(W)	75.0	74.1	1.1	23.4	90	99.6	91.1	M-high
31 F	Funk G-80	74.4	73.4	1.4	23.7	89	98.4	90.3	M-high
31 F	Hoosier Crost 840	74.4	71.5	3.7	21.9	90	99.6	87.9	Medium
33 \	Vard 120(W)	70.5	70.1	. 6	21.7	85	94.0	86.2	M-high
	Average of all entries	82.4	81.3	1.4	21.3	90.4			
		_	_						_

A difference of less than 4.1 bushels between total yields of any two entries in this summary is not significant.

			1947	RESU	LTS				
1	Stiegelmeier S-13	75.1	74.0	1.4	20.0	96	98.5	118.6	Medium
2	Producers 900	74.3	72.9	1.8	18.8	97	99.5	116.8	Medium
3	U. S. 13 (Canterbury)	74.0	72.7	1.8	19.0	96	98.5	116.5	Medium
4	Moews 830	73.6	72.1	2.0	19.2	99	101.5	115.5	Medium
5	Producers 1050	73.0	72.2	1.1	21.0	100	102.6	115.7	Medium
6	Pioneer 313B	70.7	69.9	.9	21.9	89	91.3	112.0	Medium
6	Producers 940	70.7	64.6	8.6	19.4	99	101.5	103.5	M-high
8	Morton M-380	70.6	68.9	2.6	21.1	96	98.5	110.4	Medium
9	Illinois 21 (Stone)	70.4	69.6	1.4	21.7	100	102.6	111.5	Medium
0	National 125	70.3	69.4	1.2	20.1	96	98.5	111.2	Medium
1	Ainsworth X-13-3	70.0	68.3	2.7	20.2	100	102.6	109.4	Medium
2	U. S. 13 (Stone)	69.9	68.7	1.6	19.7	98	100.5	110.1	Medium
3	Illinois 972 (Pfeifer)	69.7	67.9	2.6	21.7	100	102.6	108.8	Medium
4	Trisler T-32	69.6	67.4	2.9	20.4	95	97.4	108.0	Medium
.5	DeKalb 816	69.5	68.8	.8	19.4	95	97.4	110.3	Medium
6	Lowe 523	69.4	69.1	. 6	19.2	99	101.5	110.7	Medium
6	U. S. 13 (Mountjoy)	69.4	68.7	.8	18.7	98	100.5	110.1	Medium
8	U. S. 13 (Daily)	69.0	67.6	1.8	20.1	96	98.5	108.3	Medium
9	P.A.G. 164	68.5	66.6	2.9	19.3	100	102.6	106.7	Medium
0	Pioneer 336	68.2	67.7	1.0	19.4	99	101.5	108.5	Medium

(Table is concluded on next page)

Table 11. — SOUTH-CENTRAL ILLINOIS — concluded

		Acr	e-yield	Damaged corn in	Mois- ture in	Erect	Ratin	g for—	Compara-
Ran	k Entry		Sound	– shelled	grain at harvest	plants	Erect plants	Sound yield	height of ear
-			-	LTS —		ded	•		
21	D 4 C 172	bu. 68.1	bu. 64.0	perct.	perct. 20.3	perct.	perct. 95.4	perct. 102,6	M-high
22 22 25 25 27 28 29	P.A.G., 173. U.S. 13 (Kelly) Whisnand 905(W). U.S. 13 (Burrus) Doubet D-41. S.S. 362. U.S. 13 (Morton). U.S. 13 (Pfeifer). Kelly K-100. Illinois 201 (Burrus).	68.0 68.0 67.9 67.9 67.7 67.5 67.1	67.1 67.0 66.2 67.4 66.1 67.6 63.5 66.0	1.3 1.4 2.7 .8 2.8 .3 6.1 1.6	20.3 20.6 18.3 19.8 20.3 20.2 20.7 19.8 19.4 19.5	99 97 99 97 100 100 100 96 96	101.5 99.5 101.5 99.5 102.6 102.6 102.6 98.5 98.5	107.5 107.4 106.1 108.0 105.9 108.3 101.8 105.8	Medium Medium Medium Medium Medium Medium Medium Medium Medium
32 34 35 36 37 38 38	Ainsworth X-21. Illinois 21 (Daily) S.S. 342. Kelly K-374. DeKalb 875. Embro 36. Daily DX9 Whisnand 804 Ward 120A. Pointer Brand 87.	66.7 66.5 66.5 66.3 65.9 65.8 65.0 64.7 64.7	65.5 65.8 62.4 64.8 64.7 62.2 64.0 64.5 63.8 63.1	1.7 1.3 6.3 2.4 1.6 5.4 1.6 .4 1.7 2.4	19.3 18.0 21.2 18.9 23.4 19.7 21.3 19.5 21.3 20.1	100 100 98 94 95 99 98 98 96	102.6 102.6 100.5 96.4 97.4 101.5 100.5 100.5 98.5 99.5	105.0 105.4 100.0 103.8 103.7 99.7 102.6 103.4 102.2 101.1	Medium Medium Medium Medium Medium M-high Medium Medium
42 43 43 45 46	Keystone 38 Bear OK-40. Hoosier Crost 746 Morton M-12. Kelly K-99. Powers 149. Illinois 126 (Daily). Trisler T-22. P.A.G. 392. Morgan M-546.	64.3 63.8 63.6 63.5 63.4 63.1 63.1 63.0	63.3 63.4 63.3 63.0 61.3 62.7 62.6 62.5 62.4 61.3	1.4 .7 .4 1.2 3.4 .9 .7 1.1	18.8 18.9 19.3 19.8 20.4 18.6 20.6 21.3 19.7 20.7	98 98 100 97 99 98 97 98 99	100.5 100.5 102.6 99.5 101.5 100.5 99.5 100.5 101.5 102.6	101.4 101.6 101.4 101.0 98.2 100.5 100.3 100.2 100.0 98.2	Medium Medium Medium Medium Medium Medium Medium Medium Medium
52 53 53 55 56 57	Farmcraft FC-81. Crow 607 Canterbury 404. National 125-1. Pioneer 300. Kelly K-88. United U-50. Holmes Utility 29H Lowe 555. United U-47.	62.6 62.4 62.2 62.2 62.0 61.8 61.6 60.4 60.2	61.7 61.6 60.9 59.5 60.1 61.5 60.6 59.9 59.0 58.6	1.5 1.2 2.2 4.3 3.2 .3 1.5 .8 2.0	19.2 19.6 20.8 19.2 19.5 18.7 18.4 22.2 20.8 18.9	96 98 97 99 97 100 99 96	98.5 98.5 100.5 99.5 101.5 99.5 102.6 101.5 98.5 100.5	98.9 98.7 97.6 95.4 96.3 98.6 97.1 96.0 94.5 93.9	Medium Medium M-high Medium M-high Medium Medium Medium Medium
62 62 64 65 65 67 68	Embro 49. Lowe 560 Pioneer 332 Ainsworth X-14A Farmcraft PC-63. Keystone 45. Ward 120(W) Morton M-33. P.A.G. 612(W) DeKalb 898	58.8 58.4 58.2 57.9 57.9 57.4 56.5 56.4	58.2 58.1 57.4 56.5 56.9 57.0 56.8 55.2 54.0 55.6	1.1 .7 1.4 2.7 1.5 1.8 1.3 1.9 4.0	21.7 21.3 25.1 24.7 18.1 23.6 21.8 21.8 26.8 23.8	97 96 100 98 92 98 90 99 94	99.5 98.5 102.6 100.5 94.4 100.5 92.3 101.5 96.4 100.5	93.3 93.1 92.0 90.5 91.2 91.3 91.0 88.5 86.5	M-high Medium Medium M-high Medium M-high Medium Medium Medium
72 73 74 75 76 77 78 79	P.A.G. 170. Crow 805. Hoosier Crost F-169. Hoosier Crost 840. Crow 608. Lowe 820. Funk G-80. Hoosier Crost 404(W). Whisnand 917(W). DeKalb 923(W). Pioneer 505(W). Average of all entries.	55.9 55.8 55.3 55.1 54.6 53.3 53.2 52.8 52.5 51.7 48.1 63.8	53.7 55.3 54.4 53.5 53.4 52.3 52.4 51.8 52.2 49.7 46.3 62.4	3.7 .9 1.7 3.2 2.1 2.0 1.9 1.9 1.0 3.8 3.5	22.4 19.8 20.3 21.5 19.3 25.0 23.9 18.4 23.4 25.9 25.4 20.6	100 99 97 100 96 97 93 100 99 97	102.6 101.5 101.5 99.5 102.6 98.5 99.5 4 102.6 101.5 99.5	86.0 88.6 87.2 85.7 85.6 83.8 84.0 83.0 83.6 79.6 74.2	M-high Medium Medium Medium Medium Medium Medium Medium Medium Medium

A difference of less than 6.4 bushels between total yields of any two entries in 1947 is not significant.

^{*}Yield for 1947 is computed as average of Illinois 21 (Daily) and Illinois 21 (Stone). b Formerly Pfister. °Yield for 1947 was that of Illinois 201 (Burrus). d Formerly known as Producers 1040. °Yield for 1947 is computed as average of U. S. 13 from the following producers: Burrus, Canterbury, Daily, Kelly, Morton, Mountjoy, Pfeifer, Stone. f Yield for 1947 was that of Illinois 126 (Daily). « Yield for 1947 was that of Illinois 972A-1 (Pfeifer).

Table 12. — SOUTHERN ILLINOIS: Alhambra

		Aoro	-yield	Damaged corn in	Mois- ture in	Erect		g for—	Compara
Ran	k Entry		Sound	shelled	grain at harvest	plants		Sound yield	height of ear
	SU	MMA	RY:	1944,ª	19 4 6, ar	d 1947			
		bu.	bu.	perct.	perct.	perct.	perct.	perct.	
1	Whisnand 917(W)	52.0	50.9	. 7	24.8	66	106.4	113.9	M-high
2 ^b	Illinois 200	50.1	49.6	1.0	22.7	60	96.8	111.0	Medium
3°	U. S. 13	49.7	49.0	1.3	21.4	72	116.1	109.6	Medium
4	Funk G-80	47.7	46.9	1.6	24.2	55	88.7	104.9	Medium
5	Pioneer 332	47.3	47.0	1.5	22.4	64	103.2	105.1	Medium
6	DeKalb 816	46.0	45.5	. 7	21.5	70	112.9	101.8	M-low
7	Hoosier Crost 840	45.8	45.2	.9	21.6	45	72.6	101.1	Mediun
7	Illinois 784	45.8	45.2	1.3	25.3	59	95.2	101.1	Mediun
9	Crow 607	45.3	44.0	3.4	22.9	63	101.6	98.4	Medium
0^{d}	Pioneer 313B	44.9	44.5	1.1	22.0	66	106.4	99.6	Medium
10	Illinois 126	44.2	43.8	1.2	21.4	57	91.9	98.0	Mediun
2	Pioneer 300	43.0	42.8	.6	20.1	65	104.8	95.7	Mediun
3	Lowe 840	41.9	40.9	2.5	24.1	68	109.7	91.5	Mediun
4f	Hoosier Crost 1005A	39.3	38.8	1.3	24.6	53	85.5	86.8	Mediun
15	Hoosier Crost 746	37.3	36.9	.9	22.0	64	103.2	82.6	M-low
	Average of all entries	45.4	44.7	1.3	22.7	62.0			

A difference of less than 6.7 bushels between total yields of any two entries in this summary is not significant.

			1947	RESU	LTS				
1	Illinois 2214(W)	71.0	68.0	4.2	26.6	44	79.4	143.4	Medium
2	Illinois 1459	65.0	63.9	1.7	28.1	67	120.9	134.8	M-high
3	Doubet D-41	64.7	62.7	3.1	26.4	66	119.1	132.3	M-high
4	Whisnand 905(W)	63.9	62.9	1.6	25.5	41	74.0	132.7	Medium
5	S.S. 903(W)	62.1	61.2	1.4	24.4	54	97.5	129.1	M-high
6	U. S. 13 (Haudrich)	61.8	59.3	$\frac{4.0}{2.4}$	22.1 26.5	61	110.1	125.1	Medium Medium
7 8	Keystone 106(W) Whisnand 917(W)	61.6 59.3	60.1 58.3	1.7	24.1	67 72	120.9 130.0	$126.8 \\ 123.0$	M-high
9	Lowe 855(W)	57.5	56.1	2.4	23.2	62	111.9	118.4	M-high
10	Ainsworth X-13-3	56.4	55.2	2.4	22.1	65	117.3	116.4	Medium
11	Bear OK-50A	56.3	55.7	1.0	21.4	67	120.9	117.5	M-low
12	Pointer Brand 87	56.2	55.1	2.0	22.8	46	83.0	116.2	Medium
13	Illinois 200 (Haudrich)	56.1	55.6	.9	23.6	48	86.6	117.3	Medium
14	U. S. 13 (Kelly)	55.9	54.6	2.4	23.6	64	115.5	115.2	Medium
15	P.A.G. 173	55.3	54.8	.,9	20.9	47	84.8	115.6	Medium
16	DeKalb 923(W)	55.2	54.7	. 9	26.4	58	104.7	115.4	Medium
17	P.A.G. 612(W)	53.8	53.0	1.4	26.5	52	93.9	111.8	M-high
18	Illinois 784 (Haudrich)	53.6	52.9	1.3	28.2	59	106.5	111.6	Medium
19	Pioneer 505(W)	53.5	53.0	1.0	27.3	61	110.1	111.8	M-high
20	Huey H-23	53.4	51.8	3.0	21.2	53	95.7	109.3	Medium
21	DeKalb 875	52.4	51.5	1.7	23.7	60	108.3	108.6	M-low
22	U. S. 13 (Canterbury)	52.3	50.9	2.6	22.1	46	83.0	107.4	M-high
23	Pioneer 302	52.2	51.7	. 9	28.5	50	90.2	109.1	Medium
24	Whisnand 804	51.8	50.0	3.4	19.2	47	84.8	105.4	M-low
25	Ward 120A	51.7	50.2	2.9	23.7	48	86.6	105.9	Medium
26	Ainsworth X-14A	51.4	49.8	3.0	25.1	74	133.6	105.1	M-high
27	P.A.G. 617(W)	51.3	50.8	1.0	25.9	65	117.3	107.2	M-high
28	P.A.G. 392	50.8	50.1	1.4	19.7	71	128.2	105.7	Medium
29	Keystone 38	50.7	50.0	1.4	22.2	46	83.0	105.5	Medium
30	Kelly K-99	50.4	49.8	1.1	19.6	37	66.8	105.1	Medium
30 32	Illinois 1453	50.4	49.4 49.8	2.0	30.7 23.5	61 42	110.1 75.8	104.2 103.8	Medium Medium
33	Bear OK-315(W)	50.3 50.0	49.8	.9 1.6	20.9	65	117.3	103.8	Medium
33	Ward 125	50.0	49.2	1.8	20.9	50	90.2	103.6	Medium
35	Hoosier Crost 708(W)	49.9	48.2	3.4	24.7	46	83.0	101.7	M-high
36	Illinois 21 (Haudrich)	49.6	48.5	2.2	21.0	66	119.1	102.3	Medium
37	Keystone 45	49.3	48.6	1.3	24.9	62	111.0	102.5	Medium
38	Pioneer 313B	49.1	48.0	2.3	22.0	50	90.2	101.3	Medium
39	Illinois 972 (Pfeifer)	49.0	46.9	4.2	24.3	40	72.2	98.9	Medium
40	Illinois 200 (Burrus)	48.9	48.3	1.3	23.8	49	88.4	102.0	Medium
41	Bear OK-69	48.6	47.5	2.3	23.2	69	124.5	100.2	Medium
42	Illinois 2216(W)	48.4	47.4	2.1	25.1	55	99.3	100.0	Medium
43	Funk G-80	48.2	46.4	3.8	25.6	66	119.1	97.9	Medium
44	Kelly K-100	48.0	47.4	1.2	20.1	46	83.0	100.0	Medium

(Table is concluded on next page)

Table 12. — SOUTHERN ILLINOIS: Alhambra — concluded

		Acre	-yield	Damaged corn in	Mois- ture in	Erect	Ratin	g for—	Compara - tive
Ran	k Entry		Sound	- shelled sample	grain at harvest	plants	Erect plants	Sound yield	height of ear
		1947	RESU	JLTS —	conclu	ded			
		bu.	bu.	perct.	perct.	perct.	perci.	perct.	
45	Pioneer 300	47.9	47.5	. 8	22.0	50	90.2	100.2	Medium
45	Huey H-20	47.9	47.3	1.3	22.0	54	97.5	99.8	Mediun
17	Kelly K-374	47.7	46.8	1.8	20.5	50	90.2	98.7	Mediun
18	Pioneer 332	47.1	46.2	2.0	23.2	34	61.4	97.5	Mediun
	Lowe 820	47.1	45.6	3.2	26.5	62	111.9	96.2	Mediun
	United U-68	47.0	46.5	1.1	23.4	65	117.3	98.1	Mediun
1	Embro 36	46.9	45.7	2.6	20.6	62	111.9	96.4	Mediun
	United U-49	46.8	45.6	2.6	23.0	48	86.6	96.2	Mediun
	Illinois 126 (Canterbury)	46.8	45.5	2.7	22.9	32	57.8	96.0	Mediun
2	Hoosier Crost FD-8	46.8	45.1	3.7	19.9	51	92.1	95.2	M-low
56	Daily DX-9	46.6	45.1	3.3	22.4	57	102.9	95.2	Mediur
	Appl A-130	46.5	45.7	1.8	23.9 27.1	67	120.9	96.4	Mediur
	Hoosier Crost 1010 DeKalb 898	46.3 46.2	45.3 45.8	.9	27.1	64 62	115.5 111.9	95.6 96.6	Mediun Mediun
	Illinois 126 (Haudrich)	45.7	45.4	.6	21.6	46	83.0	95.6	Mediur
	DeKalb 816	45.5	44.8	1.6	22.5	52	93.9	94.5	M-low
	Lowe 830	45.3	44.4	2.0	27.4	36	65.0	93.7	Mediun
	Whisnand 834	44.6	42.0	5.7	25.9	44	79.4	88.6	Mediun
	P.A.G. 164.	44.2	43.0	2.7	21.5	73	131.8	90.7	Mediun
	Morgan M-546		41.7	2.5	21.2	59	106.5	88.0	M-low
55	Lowe 523	42.7	41.8	2.1	22.3	44	79.4	88.2	Mediur
	U. S. 13 (Morgan)	40.8	39.9	2.2	23.6	71	128.2	84.2	Mediur
7	Lowe 840	40.4	37.9	6.2	27.5	54	97.5	80.0	Mediur
	P.A.G. 170	39.8	39.6	.6	20.6	42	75.8	83.5	M-low
9	Embro 49	39.6	38.3	3.2	23.8	49	88.4	80.8	Mediur
0	Hoosier Crost 840	39.5	38.7	1.9	24.0	67	120.9	81.6	M-low
1	Crow 607	38.5	36.6	4.9	25.6	46	83.0	77.2	Mediur
2	S.S. 478	38.4	36.9	3.8	24.1	49	88.4	77.8	Mediu
13	National 129	37.7	37.5	. 6	20.6	48	86.6	79.1	M-low
4	Farmcraft FC-88	36.1	31.7	12.1	24.6	43	77.6	66.9	Mediu
75	Hoosier Crost 746	35.8	35.3	1.4	21.7	56	101.1	74.5	M-low
6	Pioneer 510(W)	35.2	34.3	2.6	24.7	66	119.1	72.4	Mediu
	Hoosier Crost 1005A	35.2	34.1	3.0	26.0	62	111.9	71.9	Mediur
78	Funk G-90	35.0	33.9	3.1	26.8	71	128.2	71.5	Mediu
79	Crow 608	33.2	32.1	3.3	21.2	66	119.1	67.7	Mediur
80	Brown K110(W)		30.3	1.8	27.4	74	133.6	63.9	M-high
81	National 125	27.1	23.4	13.8	23.0	43	77.6	49.4	Mediun
	Average of all entries	48.5	47.4	2.5	23.7	55.3			

A difference of less than 14.2 bushels between total yields of any two entries in 1947 is not significant.

^{*1945} data omitted because the crop did not mature. b 1947 yield was average of Illinois 200 produced by Burrus and Haudrich. c 1947 yield was average of U. S. 13 produced by Canterbury, Haudrich, Kelly, and Morgan. d Averaged with Pioneer 313D which appeared in 1944 tests. c 1947 yield was average of Illinois 126 produced by Canterbury and Haudrich. d Averaged with Hoosier Crost 1005 which appeared in 1944 tests.

Table 13.—EXTREME SOUTHERN ILLINOIS: Dixon Springs
Bottomland and Upland

548

		Damaged Acre-yield corn in		Mois- ture in	Erect	Rating for-		Compara- - tive	
Rai	ik Entry		Sound	shelled	grain at liarvest	plants	Erect plants	Sound vield	height of ear
	SUMMA	RY:	Botto	omland	1945, 19	46, an	d 1947		
		bu.	bu.	perct.	percl.	perct.	perct.	perct.	
1	Whisnand 905(W)	63.9	63.2	1.2	23.9	84	102.9	121.3	M-high
2	Keystone 106(W)	61.6	61.1	.9	23.9	85	104.2	117.3	M-high
3	Whisnand 917(W)	59.5	58.8	1.2	23.3	84	102.9	112.8	M-high
4	P.A.G.* 612(W)	58.2	57.7	. 8	24.1	82	100.5	110.7	Medium
5	Lowe 855(W)	57.4	56.0	2.8	23.1	77	94.4	107.5	M-high
6	U. S. 13 (Pfeifer)	53.0	52.2	1.4	22.0	91	111.5	100.2	Medium
7	Whisnand 834	51.7	51.3	1.0	24.8	73	89.5	98.5	Medium
8	Hoosier Crost 707(W)	51.6	51.1	1.1	26.0	79	96.8	98.1	Medium
9	Lowe 840	51.0	50.5	1.3	21.5	76	93.1	96.9	Medium
10	Morgan M-546	49.8	49.0	1.5	21.8	79	96.8	94.0	Medium
11	Ward 120A	49.0	48.7	. 6	21.2	86	105.4	93.5	Medium
12	Hoosier Crost 840	46.0	44.3	2.9	20.8	86	105.4	85.0	Medium
13	Hoosier Crost 1010	44.3	43.8	1.2	25.0	82	100.5	84.1	Medium
14	National 129	42.8	41.9	1.9	27.5	79	96.8	80.4	Medium
	Average of all entries	52.8	52.1	1.4	23.5	. 81.6			

A difference of less than 5.7 bushels between total yields of any two entries in this summary is not significant.

		ESUI	TS:	Bottomla	Bottomland				
1 1 3 4 5 6 7 8 9	P.A.G. 620(W) Ward 135(W) P.A.G. 612(W) Doubet D-41 Whisnand 917(W) Illinois 784. Ward 130 Keystone 106(W) Bear OK-315(W) Illinois 2216(W)	52.7 52.7 52.1 51.3 51.2 49.7 48.8 48.2 48.1 48.1	51.6 51.3 51.6 50.3 50.9 48.9 47.8 47.6 48.0 47.3	2.0 2.6 .9 2.0 .6 1.6 2.1 1.3 .2	28.1 33.4 29.5 26.7 26.9 28.5 33.6 26.7 24.3 27.0	74 69 79 85 80 69 56 78 74	92.0 85.8 98.2 105.7 99.5 85.8 69.6 97.0 92.0 111.9	126.2 125.4 126.2 123.0 124.4 119.6 116.9 116.4 117.4 115.6	M-high M-high Medium Medium M-high Medium M-bigh M-bigl Medium Medium
11 11 13 14 15 16 17 17 19 20	Illinois 1459. Lowe 865(W). Hoosier Crost 708(W). S.S. 903(W). Whisnand 905(W). Embro 155(W). Pioneer 505(W). United U-6(W). DeKalb 982(W). Lowe 855(W).	47.2 47.2 47.0 46.5 46.2 46.1 46.0 45.7 45.6	45.8 45.3 46.3 45.7 45.3 45.5 45.8 45.3 45.4	3.0 4.0 1.5 1.8 1.9 1.2 .5 1.6 .7	34.4 25.3 28.2 24.0 27.7 24.6 27.4 31.3 31.3 27.5	87 79 82 78 79 80 83 81 93 75	108.2 98.2 102.0 97.0 98.2 99.5 103.2 100.7 115.7 93.3	112.0 110.8 113.2 111.7 110.8 111.2 112.0 110.8 111.0 105.4	M-high Medium Medium M-high Medium Medium M-high Medium Medium
21 21 23 24 25 26 27 28 29 30	Ward 125 Ainsworth X-14A Illinois 2214(W) Keystone 45 Kelly K-88 S.S. 901(W) Pioneer 304 Bear OK-69 DeKalb 923(W) P.A.G. 173	45.4 45.3 44.3 43.7 43.6 43.1 43.0 42.4 42.2	45.2 44.4 44.5 44.2 43.2 40.5 42.9 41.9 41.2 41.5	.5 2.3 1.7 .3 1.2 7.2 .4 2.5 2.8 1.6	23.9 24.3 29.5 24.8 20.7 26.1 27.0 27.8 28.2 22.6	88 77 82 78 73 72 85 82 80 84	109.4 95.8 102.0 97.0 90.8 89.6 105.7 102.0 99.5 104.5	110.5 108.6 108.8 108.1 105.6 99.0 104.9 102.4 100.7 101.5	Medium Medium Medium Medium Medium Medium Medium Medium Medium
31 32 33 34 35 36 36	Embro 49. Ward 120A. Hoosier Crost 707(W) Bear OK-40A. Whisnand 834. Kelly K-374. Lowe 840.	42.1 41.9 41.7 41.6 40.7 40.4	41.7 41.4 41.2 39.2 39.9 40.3 40.2	1.0 1.2 1.1 5.8 2.0 .2 .5	26.5 24.3 28.8 24.7 29.1 24.9 22.8	81 86 82 84 77 80 81	100.7 107.0 102.0 104.5 95.8 99.5 100.7	101.9 101.2 100.7 95.8 97.6 98.5 98.3	Medium Medium Medium Medium M-low Medium M-low

(Table is concluded on next page)

Table 13. — EXTREME SOUTHERN ILLINOIS — concluded

				Damaged corn in	Mois- ture in Erect		Rating for—		Compara-
Ran	c Entry		Sound	shelled	grain at harvest	plants	Erect plants	Sound yield	height of ear
	1947 R	ESUL	TS:	Bottom	land —	conclu	ded		
		bu.	bu.	perct.	perct.	perct.	perct.	perci.	
36 39 40	Lowe 820	40.4 39.6 38.9	39.7 38.8 38.4	1.8 1.9 1.4	23.6 20.9 24.3	79 82 79	$98.2 \\ 102.0 \\ 98.2$	97.1 94.9 93.9	Medium M-low M-low
41 42 43 44 45 46 47 48 ^b 49 50	U. S. 13 (Pfeifer) Keystone 38. Pioneer 332. P.A.G. 164. Hoosier Crost 1005A Illinois 246-2. Stiegelmeier S-13. Illinois 448. Funk G-80. Morgan M-546.	38.4 38.0 37.6 37.4 36.3 35.9 35.7 35.5 35.4 35.2	37.9 37.6 36.2 37.0 35.4 35.5 34.3 34.9 35.2 34.6	1.2 1.1 3.7 1.1 2.6 1.1 3.9 1.7 .6	25.5 28.4 24.2 21.2 28.0 25.0 22.3 30.2 24.6 25.0	85 86 84 83 92 72 84 80 82 80	105.7 107.0 104.5 103.2 114.4 89.6 104.5 99.5 102.0 99.5	92.7 91.9 88.5 90.5 86.6 86.8 83.9 85.3 86.1 84.6	M-low Medium Medium M-low Medium M-low M-low M-low M-high M-low Medium
51 52 52 54 55 56 57 58 59 60	Doubet D-42. Pioneer 300. Lowe 830. S.S. 478. Hoosier Crost 1010. Hoosier Crost 840. DeKalb 898. P.A.G. 170. Brown K-110(W) National 129 Average of all entries.	34.8 34.6 34.6 34.4 33.8 33.0 32.8 29.8 29.6 24.6 41.7	32.7 34.3 34.2 34.2 33.5 32.9 32.5 29.3 28.0 24.2	5.9 .8 1.3 .6 1.0 .4 1.0 1.8 5.4 1.6	23.0 22.1 25.1 26.3 26.3 24.0 29.6 22.5 25.5 40.7 26.5	81 72 82 80 90 83 83 84 77 84	100.7 89.6 102.0 99.5 111.9 103.2 104.5 95.8 104.5	80.0 83.9 83.6 83.6 81.9 80.4 79.5 71.6 68.4 59.2	M-low M-low Medium Medium Melium M-low Medium M-low Modium M-low

A difference of less than 6.3 bushels between total yields of any two entries in 1947 bottomland is not significant.

		1947	RES	ULTS:	Upland	l			
1	1llinois 2216(W)	38.3	37.8	1.3	19.6	85	108.0	112.8	Medium
2	Ward 135(W)	37.5	37.1	1.2	20.6	55	69.9	110.7	Medium
3	Funk G-711	37.2	36.5	1.9	23.0	71	90.2	109.0	Medium
4	Illinois 1459	36.5	36.3	. 6	24.5	81	102.9	108.4	Medium
5	DeKalb 982(W)	36.2	35.3	2.4	20.4	86	109.3	105.4	M-low
5	Whisnand 905(W)	36.2	34.8	3.9	22.8	66	83.9	103.9	Medium
7	Keystone 106(W)	35.5	34.9	1.7	21.6	81	102.9	104.2	M-high
8	Illinois 1233-1	33.4	32.4	3.0	20.2	91	115.6	96.7	M-low
9	Illinois 1445	32.7	32.2	1.4	23.9	85	108.0	96.1	M-low
9	Hoosier Crost 707(W)	32.7	31.9	2.3	20.0	82	104.2	95.2	Medium
11	P.A.G. 612(W)	31.6	31.2	1.3	18.5	75	95.3	93.1	M-low
12	Illinois 2214(W)	31.5	31.2	1.0	20.7	70	88.9	93.1	M-low
13	Lowe 855(W)	30.6	30.4	.8	22.3	77	97.8	90.7	Medium
13	Pioneer 505(W)	30.6	30.3	1.0	24.4	83	105.5	90.4	Medium
15	Brown K-110(W)	30.0	29.8	. 8	22.0	92	116.9	89.0	Medium
	Average of all entries	34.0	33.5	1.6	21.6	78.7			

A difference of less than 5.7 bushels between total yields of any two entries in 1947 upland is not significant.

^{*} Formerly Pfister. b Three of the six plots are Illinois 1459.

SOIL ADAPTATION TEST

Hybrids were tested at Urbana for their adaptation to soils differing in fertility level. The same six single-cross and three doublecross hybrids included in 1946 were used again.

Soils. The two areas used for the tests are on the Agronomy south farm and differ in productivity as a result of long-continued use of different cropping systems. In the Southwest rotation a high state of productivity has been maintained by a systematic rotation of corn, oats, clover hay, and wheat with a red-clover catch crop. The South-Central area has been depleted of fertility by a rotation of corn, corn, corn, and soybeans. Both fields have received manure and phosphate. The predominating soil type on both fields is Sidell silt loam.

Season. Wet weather delayed the planting of these fields almost a month. The highly productive field was planted June 10 and the less productive, June 12. Lack of moisture and extreme heat during the middle of the growing season made for lower than average yields on both of these fields.

1947 results. The average yields of all hybrids in the test were lower than in any previous year since 1936. The results demonstrated the ability of certain single- and double-cross hybrids to withstand adverse conditions. Single cross $Hy \times O7$ and double cross Illinois 972-1 were the two highest yielding entries on both fields (Table 14). On the highly productive field, Illinois 972-1 was highest in yield. The three single crosses that include the four inbred lines which are used in the production of Illinois 972-1 were next in order of yield.

On the medium productive field, Hy \times O7 was highest in yield followed by Illinois 972-1. In this test all the inbreds making up Illinois 972-1, with the exception of L317, were in the single crosses ranking at or near the top in yield.

On both fertility levels, the yield of early maturing single crosses and early maturing double crosses was near the bottom. Their 1947 performance agrees in this respect with their 1946 performance. The lowest yielding entry on both fields was single cross $5120 \times \text{Hy}$. This hybrid was also the lowest yielding of the nine in 1946.

Table 14. — SOIL ADAPTATION TEST: Central Illinois, Urbana, 1947

		Total	Erect	Rating for-	
Rank	Entry	acre- yield	plants	Erect plants	Total yield
HIGHL	Y PRODUCTIVE SOII rolling phase (S200			am, sligh	itly
- Alexin		bu.	perct.	perct.	perct.

		bu.	perct.	perct.	perct.
1	Illinois 972-1	84.2	87	98	114
2	Hy × O7	83.8	92	103	114
3	Hy × 317	81.6	85	96	111
	WF9 × Hy		92	103	104
5	Illinois 246	76.2	80	90	104
6	Illinois 751	68.2	81	91	93
7	WF9 × M-14	66.7	97	109	91
	WF9 × 38-11	64.6	99	111	88
	5120 × Hy	59.9	88	99	81
	Average	73.5	89		

A difference of less than 3.5 bushels between total yields of any two of the above entries is not significant.

MEDIUM PRODUCTIVE SOIL: Mostly Sidell silt loam, slightly rolling phase (S800, South-Central rotation)

			-		
1	Hy × O7.	54.5	76	97	128
2	Illinois 972-1	45.0	67	86	106
3	WF9 × Hy		93	119	103
4	WF9 × 38-11		92	118	100
5	Illinois 246		61	78	100
6	Hy × 317	41.8	76	97	98
7	Illinois 751		80	103	95
- 8	WF9 × M14		76	97	92
9	5120 × Hy		81	104	77
	Average	42.6	78		

A difference of less than 3.7 bushels between total yields of any two of the above entries is not significant.

Lodging was less severe on the highly productive soil than on the medium productive field. Single crosses were superior to double crosses in standability. Illinois 972-1 was the best of the double crosses and WF9 \times Hy, WF9 \times 38-11, and Hy \times O7 were the best of the single crosses in percentage of erect plants.

SUMMARY

In 1947 two hundred ninety-five hybrids were tested on six fields in Illinois. As in 1946 six single-cross and three double-cross hybrids were grown at Urbana on two fields differing in productivity.

Because of wet weather, planting was late. The planting dates ranged from May 24, when the fields at Galesburg and Sullivan

were planted, to June 16, when corn went in on the bottomland field at Dixon Springs. Stands on the Woodstock, Galesburg, Sullivan, and Dixon Springs bottomland fields were good; they were variable on the Sheldon, Alhambra, and Dixon Springs upland fields. Heat and drouth were severe enough to lower yields in the northern, central, south-central, and southern portions of the state. The weighted average yield of all hybrids in the tests was 56.7 bushels an acre. This is an average of 17.2 bushels an acre, or 43 percent, above the state average yield. But it is also 17 bushels an acre less than the weighted average of all hybrids in the 1942-1946 tests.

The 1947 state average, 39.5 bushels an acre, was the lowest since 1936, a pre-hybrid year. The state average for the last five years was 50.6 bushels an acre, or 28 percent above the 1947 average.

Results of the 1947 hybrid corn tests were briefly as follows:

- 1. The Galesburg field in west north-central Illinois had the highest average yield in the state, 73 bushels an acre. On the other test fields the average yields per acre were: Sheldon, 71.1 bushels; Sullivan, 63.8 bushels; Alhambra, 48.5 bushels; Woodstock, 42.3 bushels; Dixon Springs (bottomland), 41.7 bushels; and Dixon Springs (upland), 34.0 bushels.
- 2. The five highest yielding hybrids in the three-year summaries are as follows: Northern Illinois Furr 67A, Sieben S-450, Doubet D-1, Nichols 5A, and DeKalb 609; West North-Central Pioneer 339, Pioneer 304, Doubet D-72, Kelly K-374, and Holmes Utility 39; East North-Central Pioneer 313B, Pioneer 304, Morton M-380, Frey 644, and Pioneer 332; South Central Doubet D-41, Bear OK-40, Producers 1050, Illinois 21, and Pioneer 313B; Southern Whisnand 917(W), Illinois 200, U. S. 13, Funk G-80, and Pioneer 332; and Extreme Southern Whisnand 905(W), Keystone 106(W), Whisnand 917(W), P.A.G. 612(W), and Lowe 855(W).
- 3. Lodging was most severe on the Alhambra field, where it amounted to 44.6 percent (Table 1). There was least lodging on the South-Central field at Sullivan, where only 2.5 percent of the plants were lodged.
- **4.** Corn borer damage was most severe on the northern Illinois field. Over 8 percent of the plants were broken below the

ear at the point of corn borer injury. Hybrids varied in amount of infestation, but in the three-year average the difference was not significant between the hybrids on any field.

- 5. Corn on the Galesburg field was injured by the southern corn root worm. Damage to the roots and strong wind caused considerable lodging on this field.
- **6.** Ear rot percentage was low on all the fields. Sheldon had least kernel rot and Alhambra had most. The most prevalent fungus on rot damaged kernels was *Fusarium moniliforme*. *Gibberella zeae* was present in least amount. On the whole, stalk and root rots were the most serious corn diseases in 1947.
- 7. The average yield of the nine hybrids grown at Urbana on the field having high productivity was 73.5 bushels an acre. On the less productive field, the same hybrids produced an average yield of 42.6 bushels an acre.
- 8. On the high producing field, Illinois 972-1 was first in yield and Hy \times O7 ranked second. On the medium productive field Hy \times O7 was first and Illinois 972-1 was second. This 1947 performance, together with that of former years, indicates the capacity of Illinois 972-1 and one of its component single crosses, Hy \times O7, to yield well under a wide diversity of climatic and soil conditions.

Of the nine hybrids in this special test, single cross $5120 \times \text{Hy}$, in 1947 as in 1946, gave the smallest yield.

PEDIGREES OF HYBRIDS

Following is a list of open-pedigree hybrids whose performance is shown in this bulletin.

III. 21(WF9 × 38-11) (Hy × 187-2)	III. 972-1 (O7 × L317) (WF9 × Hy)
III. 101(WF9 × M14) (W26 × 187-2)	III. 1091A (WF9 × M14) (Hy × 187-2)
III. 126 (WF9 × 38-11) (Tr × L317)	Ill. 1233-1 (WF9 \times 38-11) (940 \times R59)
III. 200 (WF9 × 38-11) (K4 × L317)	Ill. 1425 (WF9 \times Hy2) (38-11 \times R61)
III. 201(WF9 \times 38-11) (187-2 \times L317)	III. 1428 (WF9 × Hy2) (R61 × 5120B)
Ill. 206 (WF9 × 38-11) (5120 × L317)	III. $1445(K4 \times 38-11) (C1.7 \times C1.21E)$
Ill. 246-2 (WF9 × Hy) (Kr-OSF × R57)	III. $1453(K4 \times 38-11) (T8 \times C1.21E)$
111. $269(W8 \times W32)(WF9 \times Hy)$	III. $1459(K4 \times 38-11)$ (CI.21E × K201C)
III. $448(38-11 \times \text{Kys}) (\text{K4} \times \text{L317})$	Ill. 2214(W) (H21 × K64) (Ky27 × R30)
III. $751(A \times 90)$ (WF9 × Hy)	Ill. 2216(W) (H21 × CI.61) (Ky27 × K64)
III. $784(Hy \times 5120)$ (K4 × L317)	U. S. 13 (Hy × L317) (WF9 × 38-11)
III. 972(WF9 \times Hy) (ITE701 \times L317)	

CONTRIBUTORS OF SEED

Appl Hybrids	Ainsworth Seed Co. Appl's Hybrid Seed Co. Bear Hybrid Corn Co. Theo. D. Brown C. E. Canterbury Seed Co. Daily's Hybrid Corn Co. DeKalb Agricultural Assn. E. W. Doubet Ed. F. Mangelsdorf & Bro., Inc.	.St. Joseph .Decatur, Box 628 .Coulterville .Cantrall .Mattoon .DeKalb .Hanna City .1020 S. 4th St.,
Formanoft Habrida	.Farmcraft Seed Co	Outand Ind
Ferris Hybrids	Ferris Hybrids	. Princeton
Frey Hybrids	Frey Hybrid Corn Co	. Gilman
	Funk Brothers Seed Co	
Halman Habrida	Furr Hybrids	. Genoa
Husbach Hubrida	. Holmes Hybrids	. Edeistein
Huor Hybrida	L. A. Huebsch & Son	Corthogo
Hulting Hybrids	.G. E. Hulting & Son	Carthage
Unnt Unbeid	Chester A. Hunt	Morrio
Illinois Umbrida	. Ill. 21 (Burrus Seed Co., Arenzville; Da	. Mollis
minois Hybrids	Co.; Haudrich Hybrid Corn Co., Bel	
	Hybrid Seed Co., Atlanta; P. A. Stone	
	Plains)	
	Ill. 126 (C. E. Canterbury Seed Co.; Da	
	Co.; Haudrich Hybrid Corn Co., Belley	
	Ill. 200 (Burrus Seed Co., Arenzville; Hau	
	Co., Belleville; Illinois Crop Improveme	nt Assn.,ª Urbana)
	Ill. 201 (Burrus Seed Co., Arenzville)	
	Ill. 206, 246-2, 448 (Ill. Agr. Exp. Sta.)	
	Ill. 751 (Ill. Crop Improvement Assn., a U	
	Ill. 784 (Haudrich Hybrid Corn Co., Bell	eville)
	Iffl. 972 (Geo. L. Pfeifer, Arcola)	
	Ill. 972-1 (Appl's Hybrid Seed Co.; Ill. C Assn., Urbana; Robt. C. Pringle, Spar	
	Ill. 1091A (Ill. Crop Improvement Assn.,	
	Ill. 1233-1, 1425, 1428, 1445, 1453, 1459, 2	2214(W), 2216(W)
77 11 17 1 1 1	(Ill. Agr. Exp. Sta.)	C. I
Kelly Hybrids	. Kelly Seed Co	.San Jose

 $^{^{\}rm a}$ Seed supplied by the Association was obtained from samples of 1946-grown hybrids submitted for the laboratory test for certification.

Lowe Hybrids Lowe Seed Co	Aroma Park
Moews Hybrids Moews Seed Co	
Morgan Hybrids Morgan Brothers	Colve
Morton Hilbridg Don A Morton & Cons	. Gaiva
Morton HybridsRoy A. Morton & Sons	Calaghana
Munson Hybrids Carl Munson	. Galesburg
National Hybrids National Hybrid Corn Co. of Ill	
Nichols Hybrids Nichols Brothers	
Null HybridsNull Seed Farms	
P. A. G. Hybrids Pfister Assoc. Growers	
Pioneer Hybrids Pioneer Hi-Bred Corn Co. of Ill	Princeton
Pointer Brand Hybrid Moore's Seed & Farm Service	. Humboldt
Powers HybridHarlin Powers	. Brocton
Pride Hybrid Pride Hybrid Co	. Glen Haven, Wis.
Producers Hybrids Producers' Crop Imp. Assn	. Piper City
Schwenk HybridsW. T. Schwenk & Sons	. Edwards
Seeber Hybrids Seeber Bros	. Champaign
Sibley Hybrids Sibley Farms Service Corp	. Sibley
Sieben HybridsSieben Hybrids	.Geneseo, R. 1
S. S. HybridsCoop. Seed & Farm Supply Service, Inc	. Muncie
Stewart HybridFrank S. Stewart	. Princeville. R. 1
Stiegelmeier HybridsH. L. Stiegelmeier	. Normal
Trisler HybridsJ. L. Trisler	. Fairmount
United HybridsUnited Hybrid Growers Assn	.Shenandoah, Ia.
U. S. Hybrids	erbury Seed Co
Daily's Hybrid Corn Co., Haudrich	
Ill. Crop Improvement Assn., * Kelly Se	
per, Quincy, Morgan Brothers, Roy A	
Mountjoy Hybrid Seed Co., Geo. L. P	
& Son)	rener, r. m. stone
Ward HybridsMontgomery Ward & Co	619 W Chicago
ward 11 of doi: 11.1.1.1.1.1.1.1.1.1.1.1.1.1.1.1.1.1.1	Ave., Chicago
Whisnand HybridsMyron Whisnand	
Williamd Trybrids Wryton Williamd	. A II COIA

^a Seed supplied by the Association was obtained from samples of 1946-grown hybrids submitted for the laboratory test for certification.

INDEX TO ENTRIES

When a hybrid appears in the summary portion of a table, the table number in this index is printed in heavy black type. At Dixon Springs the bottomland field is indicated in this index as 13B, the upland field as 13U.

Hybrid Table	Hybrid Table
Ainsworth X-13-3	Canterbury 40411
Ainsworth X-14A	Crow 360
Ainsworth X-21	Crow 407
Ainsworth X-234, 5, 6, 8	Crow 514(W)4, 4, 5, 5
Appl A-1309, 10, 12	Crow 6076, 6, 7, 8, 9, 9, 10, 10, 11, 11, 12, 12
Appl A-136	Crow 607(W)
Appl A-2029, 10	Crow 6089, 9, 10, 10, 11, 11, 12
	Crow 633
Bear OK-204, 5	Crow 80511, 11
Bear OK-40	
Bear OK-40A13B	Daily DX9
Bear OK-506, 8, 9, 10	DeKalb 404A4, 4, 5, 5
Bear OK-50A	DeKalb 4104, 5
Bear OK-606, 8	DeKalb 422
Bear OK-6912, 13B	DeKalb 458
Bear OK-88T9, 10	DeKalb 6094, 4, 5
Bear OK-315(W)12, 13B	DeKalb 615
Brown K-110(W)12, 13B, 13U	DeKalb 628A 6, 6, 7, 8, 9, 9, 10, 10

Hybrid Table	Hybrid Table
Pybrid Table DeKalb 800A	Illinois 21
De Kalb 816 6. 6. 7. 8. 11. 11. 12. 12	Illinois 21
DeKalb 817A	Illinois 21 (Daily)
DeKalb 847	Illinois 21 (Haudrich)
DeKalb 87511, 12	Illinois 21 (Mountjoy)9, 10
DeKalb 89811, 12, 13B	Illinois 21 (Daily)
DeKalb 923(W)	Illinois 101
DeKalb 982(W)	Illinois 101 (Producers)
Doubet D-1	Illinois 126
Doubet D 42 6 6 9 12 12 P	Illinois 126 (Canterbury)
Doubet D 47 0 0 10 10	Illinois 126 (Daily)
Doubet D-72 6 6 7 8	Illinois 200
Doublet B-12, 0, 1, 0	Illinois 200 (Burrus)
Embro 36. 11, 12 Embro 49. 11, 12, 13B	Illinois 200 12 12 11 13 14 15 15 15 16 17 17 17 17 17 17 17
Embro 49	
Embro 95	Illinois 201 (Burrus),11
Embro 95	Illinois 206
	Illinois 201 (Burrus)
Farmcraft FC-40. 4, 5 Farmcraft FC-47. 6, 6, 7, 8 Farmcraft FC-69. 9, 9, 10 Farmcraft FC-81. 11, 11	Illinois 246-2
Farmcraft FC-47	Illinois 2694
Farmcraft FC-69,	Illinois 269 (Producers)
Farmcraft FC-81	Illinois 448
	Illinois 246-2
Farmcraft PC-43. 4, 5 Farmcraft PC-63. 9, 10, 11 Ferris F-A1. 6, 8	Illinois 784 (Hondrich)
Forris F A1	Illinois 072 (Pfoifor)
Ferris F-A1	Illinois 972 (Fieller)
Formin F 14 0 10	Illinois 972-1 (Appl) 0 10 10
Frey 425 4, 4, 5 Frey 644 9, 0, 10, 10 Frey 645 9, 9, 10, 10 Frey 692 9, 9, 10	Illinois 972 Frener) 1, 12
Frey 644 9 9 10 10	Illinois 972A-1 11
Frey 645	Illinois 1091A 4 4 5
Frey 692	
	Illinois 1425 9, 10
Funk G-29	Illinois 1428
Funk G-37	Illinois 1445
Funk G-29. 4, 4, 5, 5 Funk G-37 6, 6, 7, 8 Funk G-80 11, 11, 12, 12, 13B	Illinois 1453. 12 13B, 13U Illinois 1459. 12, 13B, 13U Illinois 2214(W). 12, 13B, 13U Illinois 2216(W). 12, 13B, 13U
Funk G-90. 12 Funk G-94 9, 9, 10, 10 Funk G-169 6, 6, 8 Funk G-211 9, 10	Illinois 1459
Funk G-94	Illinois 2214(W)
Funk G-169	Illinois 2216(W)
Funk G-211	
Hink (*=/11	Kelly K-779, 9, 10
Furr 33	Kelly K-889, 10, 11, 13B
Furr 33 4, 5 Furr 44A 4, 5, 5 Furr 66A 4, 5	Kelly K-99
Furr 00A	Kelly K-77. 9, 9, 10 Kelly K-88 9, 10, 11, 13B Kelly K-99. 11, 11, 12 Kelly K-100. 6, 8, 11, 12 Kelly K-374. 6, 6, 8, 9, 9, 10, 10, 11, 11, 13B Keystone 38. 9, 9, 10, 11, 11, 12, 13B Keystone 42. 6, 6, 8 Keystone 44. 4, 5 Keystone 45. 11, 12, 13B Keystone 106(W) 12, 13B, 13B, 13B, 13U
Furr 67	L'evetone 39 0 0 10 11 11 12 12B
	Keystone 42 6 6 8
Holmes Utility 29 6, 6, 8 Holmes Utility 29H 11 Holmes Utility 39 6, 6, 8	Keystone 44 4 5
Holmes Utility 29H	Keystone 45
Holmes Utility 39	Keystone 106(W)
Hoosier Crost FD-34, 5	
Hoosier Crost FD-44, 5	Lowe 4(W)4, 5
Holmes Utility 39	Lowe 6(W)
Hoosier Crost 12	Lowe 15
Hoosier Crost F-138	Lowe 24
Hoosier Crost F-140	Lowe 32
Hoosier Crost F 150 4 7 0 10	Lowe 54
Hoosier Crost E-160 6 9 0 10 11	Lowe 52
Hoosier Crost F 170	Lowe 510
Hoosier Crost 404(W)	Lowe 520 6 6 7 8 0 0 10 10
Hoosier Crost 668 6 7 8 9 9 10 10	Lowe 54 4, 5 Lowe 52 4, 5 Lowe 510 6, 8 Lowe 514 6, 8, 9, 10 Lowe 520 6, 6, 7, 8, 9, 9, 10, 10 Lowe 523 6, 8, 9, 10, 11, 12
Hoosier Crost 707(W)	
Hoosier Crost 708(W)	Lowe 560. 11 Lowe 580. 9, 10 Lowe 820. 11, 12, 13B
Hoosier Crost 746 6. 8. 11. 11. 12. 12	Lowe 580
Hoosier Crost 840 11, 11, 12, 12, 13B, 13B	Lowe 820
Hoosier Crost 1005A	
Hoosier Crost 101012, 13B, 13B	Lowe 840
Huebsch H-44 4, 5 Huebsch H-55 4, 5 Huey H-20 6, 8, 12	Lowe 855(W) 12, 13B, 13B, 13U
Huebsch H-554, 5	Lowe 865(W)13B
Huey H-20	
Huey H-23 6, 8, 12 Huey H-42 6, 8, 9, 10 Huey H-50 6, 8, 9, 10 Hulcing 101 6, 8	Moews 14
Huer H 50	Moews 15
Tuey n-50	Moews 15. 4, 5 Moews 85. 4, 5 Moews 520. 6, 8, 9, 10
Hulting 240	Moove 523
Hulting 240. 4, 5 Hulting 260. 4, 5, 6, 8 Hulting 366. 6, 8 Hulting 380. 6, 8	Moews 550 6 6 7 9 0 10
Hulting 366 6 9	Moows 830
Hulting 380	Morgan M-105 4 4 5
Hunt 60(W)	Moews 523 , 9, 10 Moews 523 , 9, 10 Moews 550 , 6, 6, 7, 8, 9, 10 Moews 830 , 11 Morgan M-105 , 4, 4, 5 Morgan M-546 , 6, 6, 7, 8, 11, 11, 12, 13B, 13B

Hybrid	Table	Hybrid Tab	ole
Morgan M-600.	6, 8	Seeber 11A	10
Morton M-12 6, 6, 8, 9, 10, Morton M-33	10.11	Sieben S-340	8
Morton M-380 6 8 9 9 10	10 11	Sieben S-450	5
Munson MX. Munson M-15.	4, 5	Sibley 777	10 11
Munson M-19.	6, 8	S.S. 362	.4
National 114-1	4 4 5	S.S. 478	
National 115A	4. 5	S.S. 903(W)	B
National 118	9. 9. 10	Stewart S-8. 6, Stiegelmeier S-9H. 4,	8
National 125	11, 12	Stiegelmeier S-9H	i B
National 129	B, 13B	Stiegelmeier S-201	. 8
Nichols 5A	4, 5, 5	Stiegelmeier S-301	8
Null N-54	6, 6, 8	Stiegelmeier S-380	10
P.A.G. 52	4.5	m 1 1 m 22	
P.A.G. 54	4, 5	Trisler T-22	
P.A.G. 60	4, 5	1113101 1-02	• •
P.A.G. 164	2, 13B	United U-6(W)13	
P.A.G. 173	2, 13B	United U-36	5
P.A.G. 270		United U-47	
P.A.G. 366A	4, 4, 5	United U-49	
P.A.G. 390	6, 8	United U-50	
P.A.G. 392	B. 13II	United U-656,	8
P.A.G. 617(W)	12	United U-68	12
P.A.G. 620(W) P.A.G. 5897. 6,	13B	U. S. 13 (Burrus)	11
Pioneer 300 6 8 0 0 10 10 11 11 12 1	12 13R	U. S. 13 (Canterbury)	12
Pioneer 302 Pioneer 304 Pioneer 313B Pioneer 313B Pioneer 332 Pioneer 332 Pioneer 332 Pioneer 332 Pioneer 342 Pioneer 343 Pioneer 342 Pioneer 343 Pioneer 343 Pioneer 343 Pioneer 343 Pioneer 344 Pioneer 345 Pion	12	U. S. 13 (Daily) U. S. 13 (Haudrich)	
Pioneer 313B 6. 8. 9. 9. 10. 11. 11. 12. 1	2. 13B	U. S. 13 (Kelly)	12
Pioneer 3329, 9, 10, 10, 11, 11, 12, 1	2, 13B	U. S. 13 (Lepper)	8
Pioneer 336	11, 11	U. S. 13 (Mountjoy)	11
Pioneer 3404,	4, 5, 5	U. S. 13 (Mountjoy)	11
Pioneer 343		U. S. 13 (Pfeifer)	11
Pioneer 344			
Pioneer 505(W)	B. 13U	Ward 1104,	
Pioneer 510(W)	12	Ward 115A	
Pioneer 4758	4, 5	Ward 115C . 0	10
Pointer Brand 87	.11, 12	Ward 120A6, 6, 8, 9, 10, 11, 11, 12, 13B, 13 Ward 120(W)11,	B
Pride D-66, 9,		Ward 125(W)	зВ
Producers 305	4.5	Ward 130	
Producers 311		Ward 135(W)	
Producers 320	4, 5, 5	Whisnand 834	B
Producers 317		Whisnand 905(W)11, 12, 13B, 13B, 13 Whisnand 917(W)11, 11, 12, 12, 13B, 13	II
Producers 709	6, 8	winishand 917(W)11, 11, 12, 12, 13B, 13	D.
Producers 709	10, 10	Single Crosses:	
Producers 940	11 1	Hy x 317	14
Producers 945 Producers 1050.	. 9, 10	Hy x O7	14
Producers 1050	.11, 11	WF9 x Hy. WF9 x M-14.	14
Schwenk S-24	6,8	WF9 x 38-11	14
Schwenk S-66.	9, 10	5120 x Hy	14







1	Ì			
-				
-				
	Ì			
and the same of				

UNIVERSITY OF ILLINOIS-URBANA

C002

Q.630.71L6B BULLETIN, URBANA 514-528 1945-48

3 0112 019529343